# MISKWABI AREA WATERSHED PLAN



# TABLE OF CONTENTS

INTRODUCTION	2
VALUES AND MISSION	4
THE WATERSHED PLAN SURVEY	4
MISSION STATEMENT	5
SUMMARY OF STEWARDSHIP ACTION PLAN	6
INTRODUCTION	6
ISSUES WITH KEY ACTIONS	7
LAKE DESCRIPTION	12
HISTORY	12
GENERAL LOCATION	17
LAKE CHARACTERISTICS	20
WATER LEVELS	26
ACCESS	28
NATURAL HERITAGE WATER QUALITY HEADWATERS VEGETATION SHORELANDS WETLANDS AND STREAMS FISH COMMUNITY WILDLIFE AND WILDLIFE HABITAT SPECIES AT RISK INVASIVE SPECIES	29 29 32 35 37 42 46 52 53

PHYSICAL ELEMENTS	54
SOILS	54
MINERALS AND AGGREGATES	54
STEEP SLOPES	54
NARROW WATER BODIES	54
SOCIAL ELEMENTS	55
SOCIAL CULTURE	55
LANDSCAPE AND AESTHETICS	55
TRANQUILITY AND NIGHT SKIES	55
RECREATIONAL ACTIVITIES	56
RECREATIONAL BOATING AND SAFETY	57
RECREATIONAL BOATING CARRYING CAPACITY	57
LAND USE	60
SUMMARY	60
RESIDENTIAL OCCUPANCY	63
SEPTIC SYSTEMS AND WASTE DISPOSAL	63
LAND USE REGULATIONS	65
OTHER REGULATIONS AND GOVERNANCE	66
GOING FORWARD	67
APPENDICES	68
REFERENCES AND BIBLIOGRAPHY	68
ACRONYMS AND ABBREVIATIONS	70
LIST OF MAPS	71
LIST OF FIGURES	71
ACKNOWLEDGEMENTS	72
INTEGRAL VISION STATEMENT	73



Long Lake Photo: Susanne James



Lost Lake Creek Photo: Michael Giza

# INTRODUCTION

The lakes involved in the Miskwabi Area Watershed Plan<sup>\*</sup>-Long, Miskwabi, Negaunee and Wenona -are unique in that they form the head waters of the Burnt River system. They are fed by wetlands, springs and streams from a surrounding landscape that has, to date, remained naturally pristine.

Outside of global climate and pollution influences there are no direct sources negatively impacting the area. The fate of the innate beauty and water quality is in the hands of the community: Municipality and County, federal and provincial agencies, property owners and visitors.

With this in mind a committee of twelve property owners, both part and full-time residents, formed a watershed plan committee in 2011 sponsored by the Miskwabi Area Community Association (MACA).

Together we established core objectives:

To determine at a grassroots level the values, ideas and concerns of all stakeholders; and to use these to fashion a watershed plan for the present day and the future.

In 2012, much effort was put in by committee members to develop a comprehensive survey; send it to all property owners; in many cases hand deliver additional copies; ask for returns; and finally, analyze the information.



Wenona Lake Photo: Dave McCulloch

The results from the 414 surveys received were very gratifying:

77% return rate for all waterfront owners and overall, including backlots, 70%.

In addition to the survey, through four years of work, the committee received verbal comments and written submissions from property owners; consulted with Municipal and County officials, provincial and federal agencies as well as other experts; held discussions at lake association meetings; and conducted two watershed plan workshops. In total, this gave the committee an accurate appreciation of issues; a broad understanding of the values of stakeholders; and thus, a comprehensive basis for the development of this watershed plan.

It is a community plan based on input from the whole community and intended for use by all to preserve the water quality, natural features, enjoyed activities and property values for today's and future generations.

\*The designation Watershed Plan was used instead of Lake Plan as four lakes are involved and they form the headwaters of the Burnt River.



Miskwabi Lake

Photo: Michael Giza



Negaunee Lake Photo: Tom Pendlebury



Miskwabi Lake Road Wetland Photo: Michael Giza

# VALUES and MISSION

# THE WATERSHED PLAN SURVEY

Below are the results of the property owners' survey which give an inclusive summary of the ideas and values of those directly connected to the lakes.

### **OWNERSHIP DEMOGRAPHICS**

- Length of ownership average 19.1 years; longest 84 years
- Seasonal/Full-time 79%/21%
- Future full-time + 17% = total 38%
- Average number in household 3.7
- Average use of property 130 days a year
- Winter use 54% of owners
- Rentals approximately 8% of owners
- Future rentals + 4% = total approximately 12%

Length of ownership and future full-time residence show a strong commitment to the lakes individually, and to Haliburton generally.

### IT'S ALL ABOUT WATER

Clearly illustrated on these two bar graphs:

- Under Natural Features, lakes were considered Very Important/Important by 100% of respondents.
- Under Issues, water quality was considered Very Important or Important by 100% of the respondents.

The importance of healthy septic systems was recognized by 96% of respondents.

### Figure1 NATURAL FEATURES



### Note: All categories scored 80%+ as VERY IMPORTANT/IMPORTANT.





### Figure 3 LAND ACTIVITIES



### Figure 4 WATER ACTIVITIES



The consensus of what is Very Important/Important was high (between 80 and 100%) for Natural Features and Issues. However, when considering Land and Water Activities the consensus of what was Very important/Important had a much wider range (20 to 90%).

People have the same values but like different activities. As such, it is important that governance and stewardship accommodate a multitude of activities while at the same time protecting the quality of the lakes.

# MISSION STATEMENT

The mission statement was written by the committee based on the common ideas and values expressed in the stakeholder survey.

# MISSION

Develop and implement a plan to preserve the water quality, environmental integrity and natural landscapes of the lakes, and in so doing, ensure the lifestyle and property values for future generations.

# SUMMARY OF STEWARDSHIP ACTION PLAN

# INTRODUCTION

Through the watershed plan development process, the people of the Miskwabi area community have clearly indicated their values as well as the lifestyles they enjoy and the lake characteristics that are essential to them.

These can be summarized as follows:

- High water quality
- Natural landscapes, forests and shorelines
- Protected wetlands and streams
- Healthy wildlife and fish populations
- Low density, non-commercial residential and cottage profile
- No additional shoreline development
- Peaceful night skies
- Variety of safe, uncrowded recreational activities
- Water levels that accommodate access and recreation as well as benefit wildlife and fish habitat
- A place for present and future generations

For the Miskwabi watershed area, and Haliburton County generally, the quality of the environment is not only important in itself, but is the underpinning of the economy. Environmental health, in particular water quality and the economy go hand in hand. It is tied to natural landscapes and clean lakes. As well, for property owners not only their lifestyles but their property values are directly tied to the state of the water and land. Our plan applauds and strongly supports Dysart et al in their recognition of the essential role of the environment for everyone in the Municipality, as expressed in its Official Plan:

"The primary objective of this Plan is to enhance and preserve those environmental qualities which contribute to the attraction of the Municipality. All development proposals will be assessed for compliance with this objective.

In particular, it is the objective of this Plan that development:

- -promote a healthy and sustainable natural environment; -protect the natural characteristics and visual aesthetics of shorelines and, wherever possible, improve and restore the natural state of shorelines and shorelands; -preserve the natural state of the shoreline as much as possible, and where the shoreline has been impaired by
- past activities, restore the shoreline's natural features including but not limited to native vegetation;
- -protect significant natural heritage features;
- -conserve significant cultural heritage resources and landscapes;
- -proceed only where any affected lakes have capacity for additional development.

All new development and the redevelopment of existing properties will be considered within the context of sound environmental planning."<sup>1</sup>

<sup>&</sup>lt;sup>1</sup> Dysart et al Official Plan, pg.6

As well, the County of Haliburton must be recognized for its approach and direction in the County of Haliburton Official Plan. Its vision is:

"A County with a co-operative environment within which the different players and population groups have worked together to achieve: a sustainable natural environment; a more stable, diversified and year-round economy based on clean, small to medium sized economic activities; residential development, both seasonal and permanent, in keeping with maintaining a small town atmosphere, environmental integrity and adequate accessible services; and a strong sense of regional identity with a more inclusive and tolerant community with all it takes to provide a high quality of life for families and individuals."<sup>2</sup>

"The County Official Plan is an indication of community values, directions and policies. The Official Plan provides policy direction to governments and individuals alike. In Haliburton County many activities and services are carried out by volunteer groups, community organizations and individuals. This Plan supports both group and individual actions and provides guidance in certain areas."<sup>3</sup>

Regarding the environment, the County Official Plan states: "Haliburton County has extensive natural areas. These areas provide the basis for a wide range of natural, economic and social functions from ecological to tourism to forestry. The natural environment forms the basis for Haliburton's way of life and its stewardship is central to this plan."<sup>4</sup>

# **ISSUES WITH KEY ACTIONS**

### 1. WATER LEVELS (Watershed Plan pg.26)

### ISSUE:

In Long (especially at the west end) and Miskwabi Lakes fluctuating water levels cause environmental issues, present navigational hazards and limit boat access.

### ACTION:

Work with the Coalition for Equitable Water Flow (CEWF) and support their initiatives with Parks Canada to develop an integrated water management system for the Trent-Severn Waterway (TSW) that includes lake-specific preferred water levels for the reservoir lakes found mainly in Haliburton.

### **ISSUE:**

Wenona Lake water levels are controlled by a natural rock sill and a beaver dam built on the sill. Over the years the high water levels resulting from the beaver dam have caused concerns. On the other hand, when the beaver dam has been altered or removed, questions have been raised about flooding, property damage and low water levels.

### ACTION:

Form a committee of Wenona Lake property owners to consult with all stakeholders, the MNR, and the Municipality to develop a procedure to effectively and equitably manage water levels.

<sup>&</sup>lt;sup>2</sup> County of Haliburton Official Plan, pg.1

<sup>&</sup>lt;sup>3</sup> Ibid, pg. 2

<sup>&</sup>lt;sup>4</sup> Ibid, pg.5

# 2. WATER QUALITY (Watershed Plan pg.29)

# ISSUE:

The stakeholder survey results confirmed that water quality is the number one priority for property owners. Strong stewardship within the community will help protect water quality.

# ACTION:

- Suggest Miskwabi Area Community Association (MACA) develop a database to store all water quality historic and future tests as well as measure water quality trends.
- Continue to work with Ministry of Environment Lake Partners Program.
- Support the Coalition of Haliburton Associations (CHA) citizen science initiative to develop a cooperative approach to comprehensive water testing by associations that will meet the required protocols of the Ministry of Environment (MOE) and Ministry of Natural Resources (MNR).
- Recommend that the Municipality of Dysart et al introduce a by-law that eliminates or restricts the use of herbicides, pesticides and fertilizers.
- 3. HEADWATERS (Watershed Plan pg.32)

# ISSUE:

Headwaters are becoming recognized as being as important as wetlands to the on-going preservation of water quality in watersheds.

# ACTION:

- Through educational information and materials, communicate the role of headwaters and the responsibility of property owners to help preserve the headwaters of the Burnt River ecosystem.
- Recommend to the Municipality and County that their Official Plans give headwaters protection similar to what environmental zones and wetlands now receive.

# 4. SHORELANDS (Watershed Plan pg.35)

# ISSUE:

Natural shorelines are considered key components in the maintenance of water quality and a lake's ability to remain healthy.

### ACTION:

- Support the County's recent by-law, Shoreline Tree Preservation, which governs the removal of trees within 30 metres of a watercourse.
- Support the shoreline inventory programme sponsored by the CHA, U-Links, Fleming College and Trent University.
- Through consultation with the Municipality take action to re-naturalize the construction site above the Rock Cut Wetland along Trapper's Trail Rd. on the north side of Miskwabi Lake.
- Work with the Municipality to establish a natural buffer zone to the Trapper's Trail Rd. causeway at the east end of Miskwabi Lake.



Harrison Point, Lake Miskwabi

Original Artwork: Gladys Gottschalk

5. WETLANDS AND STREAMS (Watershed Plan pg.37)

## ISSUE:

Wetlands in the catchment area of a lake are considered critical to the lake's water quality and general health.

# ACTION:

- Work to have the five wetlands in our watershed that scored over 80% on the County of Haliburton Natural Heritage Mapping Report assessed for Provincial Significance.
- Recommend that waterways (streams and rivers) be added to the County listing of Environmental Protection Zones and included in the Municipal By-laws with the same protection that governs wetlands.
- Work with the Municipality and County to add to the Official Plan the six wetlands that do not now appear on the Environmental Protection Zones map.
- Recommend that the current Municipal by-law governing development adjacent to wetlands be updated to disallow site alterations to wetlands and other Environmental Protection Zones.
- Recommend that the Municipality support the use of the MNR *Environmental Guidelines for Access Roads and Waterway Crossings*.

# 6. FISH COMMUNITY (Watershed Plan pg.42)

## ISSUE:

It is well recognized that lake trout populations are in peril in Haliburton. Both Long and Miskwabi Lakes are classified as cold water, lake trout lakes.

# ACTION:

- Support the recommendations of Dr. David Evans regarding the importance of the release of trout caught within restricted size range (slot sizes) and the protection of lake trout lakes (like Miskwabi) that have good habitat for natural reproduction.
- Support the CHA initiative, now with the MNR, to close the lake trout season earlier to protect breeding trout.
- Encourage a "catch and release" policy for lake trout on Miskwabi and Long lakes.
- Encourage the use of lead-free fishing equipment.
- Work with the MNR to build self-sustaining lake trout populations in Long and Miskwabi Lakes.



Miskwabi North Wetland

Photo: Larry Holden

### 7. INVASIVE SPECIES (Watershed Plan pg.53)

### ISSUE:

Invasive species can cause severe, negative alterations to natural ecosystems as well as be very troublesome to people.

### ACTION:

- Continue to work with the *Invading Species Watch* programme.
- Increase educational material and signage to prevent contamination of the lakes by boats and bait buckets.
- Encourage MACA to join the new MNR/Federation of Ontario Cottagers Association (FOCA) Early Detection & Distribution Mapping System EDDMapS programme.
- 8. TRANQUILITY & NIGHT SKIES (Watershed Plan pg.55)

### **ISSUE:**

Over 80% of survey respondents indicated that "peace and quiet" and "night skies" were "Very Important" or "Important".

### ACTION:

- Support the educational handout *Guidelines for Owners, Renters and Visitors* developed by MACA. The one-page document is available for all to use.
- Develop separate educational handouts for owners who rent and renters. These would cover the environment, septic systems, waste disposal, noise and other topics related to renting.

9. RECREATIONAL BOATING & SAFETY and RECREATIONAL CARRYING CAPACITY (Watershed Plan pg.57)

### ISSUE:

With swimming enjoyed by 98% of survey respondents, various forms of boating by 80%, faster and more powerful watercraft, and crowded weekend conditions, safety has become a growing concern. This concern is supported by a review we conducted of U.S. studies regarding recreational carrying capacity of lakes.

### ACTION:

- Develop a Boating Code of Conduct for property owners and lake users explaining the rules for safe and pleasurable operation including following the Safe Boating Guide, minimizing wake near shores and narrows, respect for neighbours, use of clean boats and trailers for environmental reasons, and a no alcohol on-board policy.
- In addition to the slow speed sign at the narrows between Long and Miskwabi Lakes, add a new sign at each of the narrows farther to the west on Long Lake.
- Recommend that the Municipality and County consider using a Recreational Carrying Capacity Model as one of the criteria when considering new or increased development on/adjacent to lakes.

### 10. SEPTIC SYSTEMS (Watershed Plan pg.63)

### ISSUE:

Malfunctioning septic systems are considered one of the most significant threats to the water quality of our lakes. This fact is all the more important for our lakes because most septic systems on the lakes, excluding those on the north shore of Miskwabi Lake, were installed over 30 years ago.

### ACTION:

- Support and work with the CHA education programme regarding pumping, inspection and maintenance of septic systems.
- Recommend that the Municipality review possible alternatives to prevent the renting of a cottage to more occupants than the septic system is designed to accommodate.
- Recommend that MACA work with the Municipality to undertake a Septic Re-Inspection Programme.

### 11. LAND USE. (Watershed Plan pg.65)

### **ISSUE:**

Non-conforming land use or development on the lands adjacent to the lakes could threaten water quality, natural landscapes, and peace and quiet. This is particularly important as the lakes are a headwater area.

### ACTION:

Work with the Municipality to reinforce all zoning and environmental regulations. Continue to ensure MACA is informed of all potential re-zoning applications and has the opportunity to engage in the re-zoning assessment process.



**Original Artwork: Tim Payment** 

# LAKE DESCRIPTION

## HISTORY

The Miskwabi watershed community is made up of the area surrounding four lakes in the Municipality of Dysart, east of the Village of Haliburton: Long, Miskwabi, Negaunee and Wenona. The area is located on Precambrian rock of the Canadian Shield which is the oldest of the earth's crust, being at least 570 million years old. In the last million years, glaciers have ripped and scraped across the landscape at least four times. A coarse grained rock known as "gneiss" was left behind. The last glacier, 15 000 years ago, created Lake Algonquin, which was made up of the basins of Lakes Erie, Huron, Michigan and parts of Superior. Our four lakes were created as the ice melted.

### **First Nation's History**

The Algonkian tribes spent their summers by lakes like ours where the fishing was good, wood was readily available for fires, and where there would be an open patch of land for growing corn and potatoes. In winters, the tribes headed further north to trap beaver and other animals. They traded their meat and furs for corn with the Hurons to the south. With the arrival of Europeans in the 17<sup>th</sup> century, beaver pelts were traded for iron axes, brass kettles and beads. During this period the Iroquois gained control over the Algonkians. As Europeans became more prominent in the late 18<sup>th</sup> century, many traditional sites on sandy beaches were submerged as loggers built dams to raise water levels so that logs could be better transported.

Lake Miskwabi's name was likely taken from the Ojibwa word, "miskwes", which meant "red" and was often used to refer to a certain kind of dogwood bark which could be smoked.

### The 1800'S

The Mississauga First Nations were not closely associated with Haliburton and on November 5, 1818, they surrendered 1.9 million acres to the Canadian Land and Immigration Company including parts of Haliburton.

In 1819, Lieutenant Catty became the first white man to have set foot in Haliburton. In 1827, Lieutenant Walpole crossed Lake Miskwabi (then known as Owl Lake), made an extensive overland portage, and paddled down the York River to the Madawaska River. In 1856, the "IBO" (Irondale, Bancroft and Ottawa) rail line opened up the area for settlement and, in 1857, James Fitzgerald surveyed Dudley Township. He wrote in 1858: "Dudley, along its western boundary, is broken. Drag Lake and its surrounding ridges are in the westerly part of the township. The easterly part is very good land and contains large tracts of excellent land. The soil is sandy loam, the timber, chiefly hardwood, with a good proportion of hemlock and pine - fully 40% of the township is good arable land and well suited for farms".<sup>5</sup>

On January 13, 1859, land in the townships of Dudley and Dysart was offered to the public at 50 cents per acre by the Canadian Commissioner of Crown Land, Philip van Koughnet. By 1869, the price of Crown Land in Dysart had tripled to \$1.50 per acre.

### Logging

In the years from 1860 to 1890 pine was the primary wood that was harvested. From 1908 to 1945, maple was the wood of choice and from 1920 to 1960, spruce and hemlock were harvested. Standard Chemical, located in Donald, owned large tracts of land in the Miskwabi area. They employed Kaufman and Edwards in the 1930's to log the area in order to make charcoal and alcohol from the hardwood. The Miskwabi mill, located near the current boat launch, apparently burned in the 1940's. There also was a gravel pit

<sup>&</sup>lt;sup>5</sup> Muskoka and Haliburton: 1615 - 1875, A Collection of Documents, U of T Press, 1963

near the airstrip. An additional mill (The King Mill) belonging to W. R. Curry was located on Long Lake. Also, a mill might have been located on Negaunee. Standard Chemical sold a large tract of land to Hasley Developments, who in turn created most of the cottage development on the four lakes that we see today.



Craig & Austin Logging Camp, Miskwabi Lake 1906<sup>6</sup> Ethel Curry is standing beside her father at the left of the front row.

### Ethel Luella Curry (1902 – 2000)



Ethel, the landscape artist, had a childhood link to the lakes. Born in a log cabin in Irondale in 1902 to parents Jessie Ellen and W. R. Curry, her family moved to Miskwabi Lake in 1906. "When I was 4, we moved to Miskwabi Lake where my father was the cook at the Craig and Austin lumber camp. Our cabin was just a tiny little thing with bunkies in it."<sup>7</sup> After working at the camp for less than a year, W.R. Curry decided he wanted a less arduous life for his young family and decided that living in Haliburton Village would be better.<sup>8</sup> Getting there would be a memorable journey. He made arrangements with the steam tugboat, which normally pulled log booms, to tow the family on a raft across Miskwabi Lake and then the length of Long Lake, nearly 5 miles. Young Ethel was put in charge of the family cow and ended up, at age 4, holding the reins of the cow with one hand and her 2year-old brother's hand with the other the whole length of the raft ride!<sup>9</sup>

After graduating from the Ontario College of Art, Ethel taught at Northern Secondary School in Toronto and painted many pictures of the Haliburton landscape which would become known as "Curry Country".

Many of the Group of Seven (Franklin Carmichael, Lauren Harris, A.J. Casson, J.E.H. MacDonald, Arthur Lismer and J.W. Beatty) worked and studied with her. She brought many of them to the Haliburton Highlands and specifically to the Miskwabi area to sketch. Her life-long friend, celebrated landscape artist Doris McCarthy, also made annual spring visits to Haliburton to paint. The photo at the right<sup>10</sup> shows Ethel (right) with Doris (left) and Noreen Masters in the summer of 1934. The Ethel Curry Gallery in Haliburton is, of course, named for her.



<sup>&</sup>lt;sup>6</sup> Popple, pg.62

<sup>&</sup>lt;sup>7</sup> ibid, pg.62

<sup>&</sup>lt;sup>8</sup> He eventually founded Curry Motors.

<sup>&</sup>lt;sup>9</sup> Popple, pg.63

<sup>&</sup>lt;sup>10</sup> Ibid, pg.158

### **Trent Severn Waterway Connection**

In 1879, the Federal Government authorized construction of the Trent Canal. The original sill dam was constructed on Long Lake in 1895 as part of the system. It was rebuilt in concrete in 1921 but as the concrete was of poor quality, it had to be rebuilt in 1939.



Long Lake Dam

Photo: Michael Giza

Depletion of forests on the Trent watershed led to fierce spring flows. From 1916 to 1950, the head water lakes (Long and Miskwabi to name two) were drained to late summer levels to ensure the lower part of the system was kept up. In the late 1950's, there was a demand for more equitable distribution of water. Various lake associations in the area agreed that the "equal percentage of storage" basis should be adopted. This policy continues today (water released from each reservoir lake be an equal percentage of all lakes). Currently, local reservoir lake associations are pushing for legislation that would allow for local lakes to determine an "optimal' draw down rate.

### The Finns on Long Lake

Through the 1930' and 40's, Finnish families bought tracts of land on Long Lake, drawn to the area by the great bass fishing. As there were no roads, they accessed their cottages by boat. The Harjula family recalled that in the 1950's they would arrive by car at the Long Lake boat launch in the dark and use a flashlight to navigate their way to their white cottage on the north shore of Long Lake.



Photo: Susanne James

### **Cottage Development**

In the early 1970's Hasley Development mapped out cottage lots and sold them from an office located at what is now the junction of Wenona Lake Road and Argillite Road. A payphone was located there – the only communication people had. Prospective buyers would be instructed on how to get to the "office" and from there they were shown the lots for sale. In those days Trapper's Trail went just as far as the "airstrip" so lots on Miskwabi were accessed by using an aluminum fishing boat, then getting out and tramping through the bush to determine which lot was for them. The Airstrip is so named since there actually was a short landing strip that developers used when flying in prospective buyers The average lot on Miskwabi originally sold for \$4200 and a down payment of \$100 was needed.

Several road names on Long Lake call back to this Finnish history. Sisu Lane

who owned 100 acres on Long Lake. The

road name was later changed to Inkinen,

several other Finnish families. Their love of the sauna is why there are several lakeside saunas on the north side

Early cottagers had their own garbage dump located on Trapper's Trail near Wenona Lake Road. It was closed by the Ministry of the Environment in the early 1980's.

<sup>&</sup>lt;sup>11</sup> Interestingly, Sisu is Finnish for determination, bravery, and resilience – much needed to access the lake in the early days.

The image below is of the sales brochure cover. Note the naming of the lakes. Some Ontario official records and maps still recognize Little Dudman as the official name of Wenona. The former name of Negaunee on MNR maps is Atwell's Lake.<sup>12</sup>



Figure 5 Developer's Sales Brochure

### The Legend of Miskwabinish

There are no facts to support the Legend, but it is said Miskwabinish, an Algonquin, made his home on what is now called Lake Miskwabi, just south of the entrance to Long Lake.

In 1784, an Algonquín war party, led by Mískwabínísh, ambushed a band of Hurons on Balsam Lake in the narrows near Cobíconk, killing all save a young maíden, Wenona, a Huron princess and a beautíful woman. Mískwabínísh took her captíve, then finding out later that she was of royal Indían blood, married her and they had three

children, only one of whom lived, a boy, named Negauree. During the period of 1782 to 1786, Miskwabinish became a powerful warrior, leading many successful raiding parties throughout Huronia to Georgian Bay and as far away as the Finger Lakes and Lake Champlain, in Iroquois territory. As a result of his strength and courage, Miskwabinish was revered by the Algonquins and feared by his enemies. In November 1796, Mískwabínísh, Wenona, and Negaunee were known to be preparing to set out by canoe for the trading post at Rosedale, for winter supplies. Their usual route would have been down Long Lake, Loon Lake and the Burnt River but other Algonquins on Long Lake, expecting to accompany them on their way, waited, then searched and found their entire camp gone. It is expected that enemy raiders surprised them, entering Miskwabi by portaging from Lakes Kennebik and Yankton and then the Burnt Creek. The mystery is complete, as all possessions, teepee, canoe, trapping supplies, food caches and dogs all disappeared and not a trace has been found to this day. Miskwabinish was thought to have a secret cave somewhere in the surrounding hills and it is believed that he was furtively followed to the cave, killed and with Wenona and Negaunee, and all their possessions, sealed therein. No one knows what happened, but Legend has it the Loon's call turned from a very happy one to very sad when Miskwabinish disappeared. To this day they say the Loon's plaintive call is the spirit of Miskwabinish lamenting his longing for the lakes he misses so much.13

<sup>&</sup>lt;sup>12</sup>References to a local family of Atwells in Wilberforce can be found online.

<sup>&</sup>lt;sup>13</sup> Attributed to Hasley Development

### Finding a Piece of the Past



Stories go around of artifacts found from First Nations or early European presence on the lakes but in the mid 1950's, as a little girl while she was skipping stones, Pam Cox found a piece of chert shaped as a spearhead at the shoreline near the Long Lake dam where her family owned land (and still does). Interestingly, chert is not local to our area and comes from Huronia, so who knows, the Legend of Miskwabinish just might have a grain of truth??? Pam speculates that she never found anything else after that because the level of the lake is never taken down as far as it was in the 1950's.

Pam's chert spearhead

Miskwabi Area Community Association formerly the Miskwabi Area Cottager's Association (MACA)



Figure 6 MACA Original logo

The Miskwabi Area Cottagers' Association was formed in 1974 at a preliminary meeting held at a cottage on Trapper's Trail Road.

The first president of the association, Jerry Strickland, set the environmental protection tone for the association that continues today. He was instrumental in uniting Long, Negaunee, Wenona and Miskwabi Lakes into one association and having representatives from all the lakes on the Executive: a practice that continues today. His environmental initiatives included ensuring that the Miskwabi Lake North Shore development happened only after all environmental aspects of the area had been carefully examined. He served as President until well into the late 80's and went on to become Executive Director of the Federation of Ontario Cottagers Association (FOCA). The "Jerry Strickland Award" is still presented each year by FOCA to worthy cottage associations for environmental projects. Both Strickland Road and the Jerry Strickland Memorial Boat Launch are named after him. Five presidents have served the association since then and although the principles remain the same, the association has grown in

membership and character. In 2013, the Miskwabi Area Cottagers' Association changed its name to Miskwabi Area Community Association (MACA) to reflect the changing profile of the property owners (cottagers and permanent residents) and amalgamation with the Miskwabi Lake North Shore Cottager's Association (MLNCA). With over 200 members, MACA continues to be a strong advocate for water quality, a well monitored environment and the local community.



Figure 7 MACA Logo adopted in 2012

# **GENERAL LOCATION**

The Miskwabi area lakes are located in the Municipality of Dysart et al, County of Haliburton, about 12 kilometres east of the Village of Haliburton via HWY. 118. Geographically speaking, Miskwabi Lake is at latitude 45°02'06" and longitude78°22' 10". The lake is located at the centrepoint between the Equator and the North Pole. The lakes are Miskwabi, Long, Negaunee and Wenona (formerly Little Dudman). Two small lakes, encircled by private lands, Hunter Lake (also known as Lost Lake) to the east and Lily Lake (local name) to the north-west, as well as a number of wetlands, drain into Miskwabi Lake.



Map 1 Topographic Map Miskwabi Area Watershed



Image of the Headwaters of the Burnt River showing the Miskwabi Area Lakes.

Map 2 Satellite Image of Miskwabi Area Watershed Credit: Google Earth

The four lakes form the headwaters of the Burnt River system which is part of the Trent-Severn Waterway. The Burnt River flows from Miskwabi down through Kinmount and into Cameron Lake at Fenelon Falls. There it joins the Trent River flowing into Lake Ontario at Trenton. To the east and northeast of Miskwabi Lake is the Irondale River and to the north and west the Drag River. These two rivers flow south to join the Burnt River north of Kinmount. On the map below the Burnt River is indicated in red flowing from the Miskwabi headwaters to Cameron Lake.



Map 3 Miskwabi Area Water Flow Map

# LAKE CHARACTERISTICS

Geologically speaking, the Miskwabi area lakes are part of the Canadian Shield located in the Algonquin Dome portion of the Laurentian Highlands.



Map 4 Physiographic Map of Southern Ontario Credit: MNR

The bedrock is granite gneiss often found as steep cliffs or outcroppings along the shorelines. The soil substrates are generally thin and shallow with some heavier glacial gravel deposits. The drainage basin is steep to gently rolling with a maximum elevation of approximately 80 metres (262.5 ft) above the lake levels. The highest land elevation above sea level in the area is about 442 metres (1450 ft) and this compares to the highest elevation in Haliburton County at 563 metres (1847 ft).

The lake surface elevation of Negaunee Lake is 404 metres (1325 ft) above sea level and close to the highest in the Haliburton Highlands.

The geological age of the granite gneiss bedrock in the watershed is 570 – 1600 million years, Late to Middle Precambrian. The colour-coded and numbered map following indicates the types of rocks in the Miskwabi area:

- White (9) Conglomerate, greywacke, arkose, calcareous sandstone and siltstone, shale, and derived metamorphic rocks
- Blue (12) Marble, calc-silicate rocks, skarn
- Rose (13) Anorthositic to tonalitic rocks



Map 5 Map 2441, Ontario Geological Survey. Credit MNR

Due to the makeup of the surrounding geology the lakes are characterized by soft water, a neutral PH, good calcium levels and are not threatened by acid rain.

# What type of Lakes do we live by?

Three basic types of lakes are found in Ontario.

### Oligotrophic Lakes

- Generally deep
- Minimal aquatic plant growth
- Low nutrient levels
- Support cold-water fish such as lake trout and whitefish
- Low levels of phosphorus and chlorophyll
- Total phosphorus test indicates < 10ug/L</li>

### Mesotrophic Lakes

- Medium depth
- Usually good for fishing; supporting a wide variety of fish such as pickerel and bass
- More nutrients than oligotrophic lakes, but not nearly as much as eutrophic lakes
- Occasional algae bloom at the surface
- Total phosphorus test indicates 10 20 ug/L

### **Eutrophic Lakes**

- Generally shallow with abundant vegetation
- Support warm-water fish such as perch, bass and pan fish
- Frequent algae blooms
- Susceptible to oxygen depletion
- High phosphorus or chlorophyll readings
- Total phosphorus test indicates >20 ug/L

Eutrophication is a lake's aging process. Sediments, erosion and the growth and decomposition of aquatic plants eventually fill up the lake bottom. Over time, the lake is converted to a wetland (e.g., a bog or marsh) and later, dry land. This process normally takes tens of thousands of years, but human activity can accelerate lake eutrophication by contributing excessive nutrients.







Long, Miskwabi and Wenona Lakes are oligotrophic lakes. Negaunee Lake, moderately enriched with good depth and temperature stratification, is on the border of oligotrophic and mesotrophic classifications.<sup>14</sup>

<sup>&</sup>lt;sup>14</sup> A Shoreline Owner's Guide to Lakeland Living, pg.2-3

# Bathymetry Maps of the Four Lakes Credit: MNR









n.b. Due to scales changes map dimensions are not comparable

# WATER LEVELS

The water levels of Negaunee and Wenona Lakes are controlled by natural rock sills. On Wenona Lake a beaver dam built on the natural sill has caused issues. with water levels, some flooding and property damage. Long and Miskwabi Lakes are controlled by the Long Lake dam of TSW. This dam controls a drainage area of 20.2 sq. km. and has a storage capacity volume of 664 ha/m. The maximum depth is 1.98m above the natural sill. As seen in the chart at the right seasonal water-level fluctuations can vary substantially: on average, 1.1 metres. Properly controlled water levels are important for a number of reasons:

- Boat navigation and safety
- Water access
- Removing boats and docks from lakes
- Water intake for property use
- Lake Trout spawning
- Loon and other water bird nesting

• Reptile and amphibian hibernation The Coalition for Equitable Water Flow (CEWF), representing the Trent-Severn reservoir lakes, has been working with the Ontario Waterways (Parks Canada) for a number of years. A local survey indicated that property owners strongly support the effort of the CEWF to achieve an integrated water management system with lake-specific preferred water levels.



#### Figure 8 Miskwabi – Long Lakes Water Levels

Credit: CEWF

ISSUE: In Long (especially at the west end) and Miskwabi Lakes fluctuating water levels cause environmental issues, present navigational hazards and limit boat access.

ACTION: Work with the Coalition for Equitable Water Flow (CEWF) and support their initiatives with Parks Canada to develop an integrated water management system for the Trent-Severn Waterway (TSW) that includes lake-specific preferred water levels for the reservoir lakes found mainly in Haliburton.

ISSUE: Wenona Lake water levels are controlled by a natural rock sill and a beaver dam built on the sill. Over the years the high water levels resulting from the beaver dam have caused concerns. On the other hand, when the dam has been altered or removed, questions have been raised about flooding, property damage and low water levels.

ACTION: Form a committee of Wenona Lake property owners to consult with all stakeholders, the MNR, and the Municipality to develop a procedure to effectively and equitably manage water levels.

Figure 9 Miskwabi – Long Lakes Preferred Water Levels					
Key Lake and Dam Statistics:					
Drainage area:         Lake area:         Full control level of dam:         Target full level :         Sill or deduction:         Maximum storage depth:         Maximum storage volume:         Log number & dimension:         Normal winter-set:	20.2 sq. km. 335 ha 2.29 m 100% 0.31 m 1.98 m 664 ha-m 7 @ 0.305 m plus 1 metal "half-log" @ 0.15 m 5 logs				
Water level fluctuation:	0.76 m				
Most significant Impacts of fluctuating Water Levels:					
<ul> <li><u>Water levels "too high":</u></li> <li>shoreline erosion</li> </ul>	Water levels "too low":• difficulty navigating between Long and Miskwabi Lakes• unmarked navigational hazards created• difficulties in removing boats and docks from lake• difficulty in accessing some bays in the lakes• water intake problems				
<ul> <li><u>Lake levels rising in June:</u></li> <li>breeding difficulties for aquatic birds</li> </ul>	<ul> <li>Lake levels falling in October:</li> <li>possible negative effect on lake trout spawning beds</li> </ul>				
Upper preferred water level limit: 7 logs to minimize shoreline erosion (2.29 m)	Lower preferred water level limit: 5 logs (1.53 m) to maintain adequate water levels for safe navigation				
Miskwabi Area Community Association supports a range between 2.29 m (7 logs) and 1.53 m (5 logs) during the navigational season.					
Winter-set Level: Miskwabi Area Community Association understands that the Long Lake dam traditional winter set level is 5 logs (1.53m).					
Source: MACA survey of property owners on Miskwabi and Long Lakes January 2012					

# ACCESS

All lakes can be accessed on a year-round basis from Hwy. 118 using Trapper's Trail Rd. As well, Long Lake can be accessed from Hwy. 118 via Kennaway Road.



# NATURAL HERITAGE

# WATER QUALITY

The response to our lake survey question regarding water quality was unanimous; every one of the 414 respondents, 100%, said water quality was "Very Important" or "Important". "It is all about the water" – aesthetically, recreationally, and economically. The lakes of the Miskwabi area have been tested over many years, some more than others. Our lakes have participated in the Ministry of Environment (MOE) Lake Partners Program. As well, the MOE and Ministry of Natural Resources (MNR) have conducted tests that we have been allowed to review. In addition, both Long and Miskwabi Lakes have been extensively examined by the MOE Lakeshore Capacity Assessment. Results for some of the more common tests are outlined in the table on page 29. What follows is an explanation of each test.

# Water Clarity – Secchi<sup>15</sup> disk readings

The visual clarity of the water in a lake is measured by lowering a Secchi disk in the water to the depth at which it disappears. Changes in the clarity are a useful tool in tracking the nutrient status of a lake along with other tests.

### Total Phosphorus (TP)

These readings are used to interpret the nutrient status of a lake. Increases in TP cause algae growth, decrease water clarity and, in extreme cases, lead to algae blooms. These can affect the aesthetics of a lake as well as cause taste/odour problems, and in rare cases, be a serious health hazard.

Lakes are placed in three categories with respect to trophic (nutrient) status: those with less than 10ug/L TP are considered

oligotrophic and characteristically are clear, deep and unproductive; lakes with TP between 10 and 20 ug/L are called mesotrophic and can be clear and unproductive at the lower end of the scale or susceptible to moderate algal blooms at the upper readings; and lakes over 20 ug/L PH are classified as eutrophic and may be subject to persistent, nuisance algal blooms.

#### **Dissolved Oxygen**

Dissolved oxygen is critical to life in a lake. During the summer and early fall oxygen levels in water below the thermocline (about 7 m) impact the survival of lake trout. The larger the volume of deep water with greater than 6 mg/L of oxygen, the better for lake trout.

### рΗ

This is a method of measuring the acidic levels of the water. Since acids release ions, the acid content is determined by measuring the amount of hydrogen ions. The pH scale ranges from 0 (extremely acidic) to 14 (extremely basic/alkaline) with 7.0 being neutral. Most aquatic plants and animals can survive in ranges 6.5 to 8.0.

#### Alkalinity

Alkalinity testing measures the acid-neutralizing (buffering) capacity of a lake by determining its carbonate-bicarbonate concentration. Many Haliburton lakes have low buffering capacity and are sensitive to acid rain. Lakes with less than 10mg/L may be subject to acidification.

#### Calcium (Ca)

In aquatic systems calcium is necessary for growth of crustaceans as they shed and rebuild their exoskeletons. Recent studies in Haliburton and Muskoka have documented declines in calcium in area lakes. Zooplankton, like Daphnia (native water fleas), are an integral part of the ecosystem and provide food for fish. Declines in calcium, and subsequently plankton, can lead to negative impacts on the lake food webs. Daphnia species may be jeopardized by lake calcium levels below 1.5 mg/L.

<sup>&</sup>lt;sup>15</sup> Named for its creator Angelo Secchi.

Figure 10 WATER QUALITY PARAMETERS						
PARAMETER	LONG	MISKWABI	NEGAUNEE	WENONA		
SECCHI Depth (m)* >5 oligotrophic 3-5 mesotrophic <3 eutrophic	Average over 2 years 2003 & 2005 4.25	Average over 15 years 1998 – 2012 8.5 No trend over 15 yrs.	Average over 5 years 2000 – 2004 4.82 No trend over 5 yrs.	Average of 11 years from 1992 – 2012 5.1 No trend over 11 yrs.		
Total Phosphorus (ug/L)* <10 ug/L oligotrophic 10-20 ug/L mesotrophic >20 ug/L eutrophic	Average over 2 years 2006 & 2008 6.95	Average over 11 years 2002 – 2012 5.7 No trend over 12 yrs.	Average over 2 years 2002 & 2004 10.5	Average over 8 years from 2002 – 2012 8.85		
Dissolved Oxygen (mg/L)** >6 excellent for lake trout	Average over 5 years 09/86, 08/89, 09/94, 08/99, 09/07 10 metres 2.4 20 metres .79 Trend to lower D.O. at both depths	Average over 4 years 09/75, 10/86, 09/01, 09/07 10 metres 8.7 20 metres 6.9 40 metres 4.1 Trend to lower DO at 40 m.	Average over 2 years 09/83 and 08/89 10 metres 1.0	Average over 3 years 08/79, 08/83, 08/84 10 metres 3.4 20 metres .1		
<b>pH**</b> >6.5 excellent 5.5 – 6.5 good <5.5 poor	Average over 2 years 1970 & 2007 7.5	Average over 8 years 1970,75,95,96,97,98,2001,07 7.3	Tested 09/83 8.0	Average over 2 years 08/83 & 08/84 7.9		
Alkalinity (mg/L)** <10 sensitive	Average over 2 years 33.9	Average over 8 years 32.6	Tested 09/83 47.9	Average over 2 years 08/83 & 08/84 37.6		
Calcium (mg/L)** >1.5 mg/L necessary	Recent testing 13.3	Recent testing 13.9	Recent Testing 21.4	Recent Testing 13.9		
*MOE Lake Partner Program **Tests completed by MOE or MNR personnel						

### Water quality conclusions

With regard to pH, Alkalinity and Calcium, the results for all four lakes are very positive.

Water Clarity and Total Phosphorus tests indicate that Negaunee is tending toward being mesotrophic; whereas Long, Miskwabi and Wenona are oligotrophic.

Dissolved Oxygen figures signify that Long, Negaunee and Wenona are not suitable habitat for lake trout. This is of concern for Long which has traditionally supported a lake trout population. Dissolved Oxygen tests for Miskwabi vary substantially from test-to-test years but generally are acceptable above 30 m. The bottom reading (40 m) has been trending downward. Even though the TP tests are good with no negative trends, the lower oxygen levels at depth along with the depleted oxygen levels in Long Lake give cause for concern.

### Lake capacity assessment

The goal of lake capacity assessment is to ensure sustainable development of our inland recreational lakes using a watershedbased approach. The Lake Capacity Model (LCM) <sup>16</sup>was first developed in the early 1970's to address the relationship between shoreline development and nutrient loading. It has been refined and updated, most notably with a component that links shoreline development and dissolved oxygen in lake trout lakes <sup>17</sup> "For cold-water lake trout lakes, the MNR criterion is a volume-weighted

hypolimnetic mean of 7 mg/L dissolved oxygen. Lakes are deemed to be at capacity or highly sensitive if:

i) Measured volume-weighted mean hypolimnetic dissolved oxygen, taken at the end of the summer, is consistently below the 7 mg/L criterion.
ii) Model predictions conclude that the loading associated with the existing vacant lots of record and/or new development proposal(s) will reduce the volume-weighted mean hypolimnetic dissolved oxygen below 7 mg/L".<sup>18</sup> With the opening of Miskwabi Lake to cottage development in the late 1960's and early 70's, both Miskwabi and Long Lakes were assessed under the Lake Capacity Model and deemed at capacity. Subsequently, the north shore of Miskwabi was opened to development with special covenants involving increased lot sizes, deeper setbacks and regulations regarding the shoreline buffer area. In the Dysart et al Official Plan both lakes are considered at capacity.

It has been noted that lake capacity assessment only analyzes lakes as defined above. Other impacts such as forestry practices; agriculture; and issues of social importance like boat traffic and safety, loss of wilderness habitat, noise, natural landscapes, shoreline crowding, and density issues are not covered. These have to be addressed by other planning and regulatory mechanisms.<sup>19</sup>

ISSUE: The stakeholder survey results confirmed that water quality is the number one priority for property owners. Strong community stewardship will help protect water quality. ACTION:

- Suggest MACA develop a database to store all water quality historic and future tests as well as measure water quality trends.
- Continue to work with MOE Lake Partners Program.
- Support the Coalition of Haliburton Associations (CHA) citizen science initiative to develop a cooperative approach to comprehensive water testing by associations that will meet the required protocols of MOE and MNR.
- Recommend that the Municipality of Dysart et al introduce a by-law that eliminates or restricts the use of herbicides, pesticides and fertilizers.

<sup>&</sup>lt;sup>16</sup> Lakeshore Capacity Assessment Handbook

<sup>&</sup>lt;sup>17</sup> Water Quality and Management of Lake Trout Lakes, pg.18

<sup>&</sup>lt;sup>19</sup> Water Quality & Management of Lake Trout Lakes, pg.18

# **HEADWATERS**

The catchment area of our four lakes is at the headwaters of the Burnt River. On the one hand, being at the headwaters our lakes are not affected by any upstream sources; on the other hand, these lakes are the source of water that flows into the downstream waterways.



Miskwabi Causeway Creek

Photo: Michael Giza

It is very important that from a stewardship point of view we do everything possible to protect the headwaters ecosystem and the guality of water that flows out of our area.

The essential importance of headwaters is just starting to be realized and studied. To quote the Ontario Headwaters Institute: "The health of our headwaters and their catchment areas are fundamental to the ecological integrity of Ontario. Among other benefits, headwaters and their catchment areas:

 comprise the majority of the surface area of most watersheds;

- constitute the majority of stream length in most watersheds;
- contribute the majority of the flow to most watercourses;
- provide significant portions of a stream's nutrients, organic material and sediment;
- help regulate the flow of water into, and can mitigate flooding in, a watercourse;
- furnish key habitat for breeding, feeding and sheltering upstream species, therefore nurturing the majority, and in many ways, the base, of a watershed's biodiversity."<sup>20</sup>

Poorly understood, in particular, are the innumerable small, temporary streams, swales and wet areas. These are especially vulnerable to impacts from crossings, channelization, flow diversion, grade lowering and realignment.

The Toronto Regional Conservation Authority has been studying headwaters and states "there is a growing body of evidence to suggest that headwaters are important sources of food, sediment, nutrients and flow to downstream aquatic systems...they provide water quality, storage and attenuation functions."<sup>21</sup>

ISSUE: Headwaters are becoming recognized as being as important as wetlands to the on-going preservation of water quality in watersheds.

### ACTION:

- Through educational information and materials, communicate the role of headwaters to property owners and their responsibility to help preserve the headwaters of the Burnt River ecosystem.
- Recommend to the Municipality and County that their Official Plans give headwaters protection similar to what environmental zones and wetlands now receive.

<sup>&</sup>lt;sup>20</sup> Ontario Headwaters Institute, pg. 1

<sup>&</sup>lt;sup>21</sup> Headwater Study, pg. 2

# VEGETATION

The landscape around the lakes at the head of the Burnt River is representative of the Great Lakes - St. Lawrence Forest and is now covered with mature forest. Logging at the turn of the twentieth century eliminated all the old growth and as the topography was not suitable for farming regeneration naturally occurred. There are four forest regions in Ontario: Deciduous Forest Region, Great Lakes-St. Lawrence Forest, Boreal Forest and Hudson Bay Lowlands. The Haliburton Highlands are part of the Great Lakes – St. Lawrence Forest. It is the transitional zone between the deciduous forest of south-western Ontario and the predominately coniferous boreal forest north. Throughout this region coniferous trees such as Eastern White Pine, Red Spruce, Eastern Hemlock and White Cedar commonly mix with deciduous broad-leaved species such as



Autumn on Negaunee Lake

Sugar and Red Maple, Basswood and Red Oak. Species more common in the boreal forest such as White and Black Spruce, Jack Pine, Aspen and White Birch also grow here. As well, this transitional forest region contains many species of fungi, ferns, mosses, shrubs and wildflowers.



Summer on Negaunee Lake

Photo: Tom Pendlebury

The forests around the Miskwabi area lakes are mixed with hardwood species dominating overall. The buffer zones along the lakes are made up of *Eastern Cedar*, *Balsam Fir*, *Eastern Hemlock*, White Spruce, Paper Birch, Red Maple, Black Ash, American Basswood and Trembling Aspen.

In the more well-drained uplands Sugar Maple, American Beech, Ironwood, American Basswood, White Pine, White Spruce, Eastern Hemlock, some Black Cherry, Yellow Birch and Red Oak are found.

On the forest floor, before the deciduous canopy spreads, and especially along south-facing hills, a wide range of early blooming plants brighten the spring ground. These plants need to get their energy from the sun before being shaded through the summer. Some common flowering plants in the Miskwabi area are:

### Dutchman's Breeches (Poppy Family)



Eastern Canada and U.S. in deciduous woods

Early spring flowering in April-May. Flowers are pollinated by bumblebees whose proboscises are long enough to tap the nectar.

Large-flowered Trillium



Large, solitary, waxy-white flower that turns pink with age. Has an erect stalk above a whorl of 3 broad leaves. Habitat is rich deciduous forest areas. Native North Americans used the rootstalks for medicinal purposes. Blooms in April and May.

### Spring Beauty



Low plant with loose clusters of pink/whitish flowers with a dark pink stripe. Spring perennial that is seen in patches and grows from underground tubes like a potato. Native North Americans ate the chestnut-tasting tubes.

#### Coltsfoot



### Trout Lily



### Purple Trillium



Solitary, nodding flowers rising on a stalk above 3 diamondshaped leaves. Found in deciduous woods and blooms

Source: The Audubon Society Field Guide to North American Flowers Photos: Virginia Vranckx

from April to May.

outside, nodding flower. One of the most common spring wildflowers, it can be found in large colonies from April to June. The name refers to the suggested similarity of the leaf markings to those on a trout.

Low dandelion-like plant with

solitary yellow head on a scaly

stalk. Grows in open areas like

to the suggested resemblance of the leaf to a colt's foot. The

extract of fresh leaves can be

dried for tea.

used in making cough drops or

Brownish-mottled leaves sheath the base of a stalk bearing a solitary, yellow inside/bronze

roadsides and blooms from April to June. Common name refers

# **SHORELANDS**



Photo: Andy Mosher

The shorelands of a lake are an essential component in the maintenance of water quality and the lake's ability to remain healthy. They are critical in providing habitat for many of the lake's life forms during some form of their life cycles. As well, shorelands can act as buffer zones to filter out nutrients such as phosphorus before they enter the lakes and cause weed growth, algae blooms, oxygen depletion and stressed fish populations.

The natural shoreline can be divided into three zones; beginning underwater and extending upland.

The *littoral zone* is the area from the water's edge to roughly where sunlight no longer penetrates to the lake bottom. As much as 90% of the species in the lake either pass through or live in this zone.

The *riparian zone* is the transitional land that forms a buffer along the water's edge.

The *upland zone* is the higher, drier ground adjacent to the riparian zone and is typically forested with the kinds of trees that take advantage of better drainage like Poplar, Spruce and White Birch. The natural vegetation of the riparian and uplands zones help to stabilize the soils, reduce erosion and act as a barrier to help filter out "undesirables" before they enter the lake. These can include natural runoff, septic tank seepage, fertilizers, pesticides, pet deposits and oil or gas spills. One of the main problems is phosphorus which occurs naturally and also in human-made products and wastes. Naturally-occurring phosphorus helps to nourish life in the lake, but when added to the man-made load, phosphorus over-feeds the lake causing excessive plant growth and algae blooms that consume the water's oxygen, resulting in poor water quality.



Figure 11 Shoreland Zones

Credit: The Shore Primer, pg.7
### Shoreline Tree Preservation By-Law

In 2012, the County of Haliburton, recognizing the importance of natural shoreline buffers, passed the Shoreline Tree Preservation By-Law (Preservation By-law No. 3505). This by-law applies to all lands within 30 metres of lakes, rivers and navigable waters. Its purpose is to:

- minimize the destruction or injury of trees
- regulate the removal, maintenance and protection of trees
- sustain a healthy natural environment by maintaining and improving the ecosystem services provided by trees
- maintain water quality
- prevent soil erosion and reduce airborne pollution
- protect fish spawning beds

The new by-law also encourages the rehabilitation and naturalization of shorelines which, over the years, have been altered.

Our stakeholder survey indicated that natural shorelines are Very Important/Important to 97 % of respondents, and as they are essential for the preservation of the water quality and related natural environment, this report strongly supports the by-law.



#### Miskwabi Shoreline Photo: Ken Morrison

# Shoreline Inventories

Monitoring and renaturalizing lake shorelines is one of the two most important actions shoreline property owners can take to protect water quality (the other being maintaining septic system health). During 2014-15 the four Miskwabi area lakes are being offered the opportunity to participate in a shoreline inventory programme sponsored by the CHA, U-Links, Fleming College, and Trent University.

Trained students from environmental studies and geography, using GIS mapping, will study each lake. Based on a Wisconsin model and a categorized rating system developed for Kennisis Lake, all the shorelines will be analyzed.

ISSUE: Natural shorelines are considered key components in the maintenance of water quality and a lake's ability to remain healthy.

#### ACTION:

- Support the County's recent by-law, Shoreline Tree Preservation, which governs the removal of trees within 30 metres of a watercourse.
- Support the shoreline inventory programme sponsored by the CHA, U-Links, Fleming College and Trent University.
- Through consultation with the Municipality, take action to re-naturalize the construction site above the Rock Cut Wetland along Trapper's Trail Rd. on the north side of Miskwabi Lake.
- Work with the Municipality to establish a natural buffer zone to the Trapper's Trail Rd. causeway at the east end of Miskwabi Lake.

### WETLANDS AND STREAMS

Four main area lakes and over 20 wetlands and streams within the boundaries as illustrated on the map below.



Map 11 Miskwabi Area Watershed Plan Catchment Area

As part of the head waters for the Burnt River watershed, our concerns about our natural habitat are slightly different than many other lakes as we do not receive flows from other water bodies. Being head water does carry a serious responsibility: for what happens in our water affects all other water downstream in the system. Wetlands and streams provide substantial ecological, social and economic value to the lakes by maintaining and improving water quality. Retention of rain and runoff assists in flood control and provides habitat for wildlife, nesting areas for birds, spawning sites for fish, nesting grounds for turtles, and homes and nurseries for various other species, as well as conditions that support a wide variety of vegetation.

#### Wetlands

These are areas where land is wet either seasonally or permanently. The result is water-logged soils and growth of water-tolerant trees and plants. Wetlands occur along edges of lakes, rivers and streams where water pools for at least part of the year.

Four classes of wetlands are recognized by MNR: swamps (wooded wetland), marshes (typically consisting of non-woody plants such as cat tails, sedges and reeds), fens, and bogs (both wetlands that accumulate peat and support acidic loving trees and plants).<sup>22</sup> In the watershed area we find swamps and marshes, possibly fens, but not bogs which are usually found in Northern Ontario in thinner soils. The wetland classification system recognizes that hydrological processes, dictated by climate and landscape factors, largely determine wetland form.<sup>23</sup>

Four different types of wetlands are recognized by MNR: lacustrine (associated with lakes); riverine (associated with rivers); palustrine (occurring upslope from previously mentioned types); and, isolated (receiving nutrients from precipitation, overland flow and groundwater)<sup>24</sup> All types are present in the watershed. Wetlands play a vital role in maintaining high quality water supplies by collecting rainfall, snowmelt and runoff; storing it; and then discharging it as surface runoff or groundwater flow. They act as buffer zones for various types of nutrients and pollutants. Other vital roles include providing wildlife corridors for seasonal breeding, migrations and habitat for many species.

<sup>&</sup>lt;sup>22</sup> What are Wetlands? Pg. 1-2

<sup>&</sup>lt;sup>23</sup> National Wetlands Working Group, 1997.

<sup>&</sup>lt;sup>24</sup> MNR, Biodiversity (website)

### Wetland Mapping

In the catchment area of the lakes we have identified 14 wetlands and in the Dysart et al Official Plan it appears that6 wetlands are not designated as Environmental Protection Zones. As well, some of the wetlands are possibly complex ones. Under the Ontario Wetlands Evaluation system, wetland complexes are ones where the topography of the landscape, the short distances between some wetlands and the density of wetlands per unit of areal landscape may be so complex that delineation of the wetlands into individually recognized units would not be an ecologically or functionally sound process.



Miskwabi North Wetland

Photo: Tim Payment

In the Haliburton Highland Land Trust report, County of Haliburton Natural Heritage Mapping, five of the wetlands in the Miskwabi watershed area are part of those in the County of Haliburton with the highest potential for provincial significance. In the study, Haliburton County wetlands were ranked between 1 and 100 with those ranking between 80 and 100 considered having the highest potential for being Provincially Significant Wetlands (PSW).<sup>25</sup> To make identification simpler, we have named the 14 wetlands and noted their score on the above-mentioned study. The map number refers to the wetlands map on the following page.

Miskwabi Ar	ea Wetlands	;
NAME	Score	Map No.
South Negaunee Lake	90 -99	1
Miskwabi East Causeway	80 -89	2
Miskwabi North	80 -89	3
Miskwabi Rock cut	80 -89	4
Miskwabi Downdraft	Under 80	5
Lily Lake	Under 80	6
Miskwabi Lake Road	Under 80	7
Inkinnen Trail	Under 80	8
South Wenona Lake	Under 80	9
West Wenona Lake	Under 80	10
Wenona Lake Rd	Under 80	11
Lost Lake East	80 - 89	12
Lost Lake North	Under 80	13
Burnt River	Under 80	14

In any case, PSW or not, the wetlands surrounding the Miskwabi area catchment are essential for the health of the lakes and water quality.

<sup>&</sup>lt;sup>25</sup> PSWs are those areas identified by the Province as being the most valuable. They are determined by a science-based ranking system known as Ontario Wetlands Evaluation System (OWES). Source: MNR



Map 12 Miskwabi Area Wetlands

Credit: County of Haliburton GIS & Mapping Site

#### Streams

Streams are channels of water that are either intermittent or permanent in nature and can include deltas and flood plain areas. Streams provide transportation of surface water, connect chains of lakes, help control erosion, silting and assist in flood abatement. They provide spawning, nursery, rearing, seasonal and adult habitat to a broad range of fish, turtles, other amphibians, reptiles and wildlife.



Miskwabi North Creek

Photo: Larry Holden

Intermittent streams control water in peak seasons and contain structure to help control channeling and silt abatement. Permanent steams flow year round and can be connecting channels between bodies of water. These streams can be either warm or cold water. In most cases cold water streams are, to a certain extent, spring fed. Consistent with past usage and policy within MNR, the definition of a coldwater stream is *one supporting or capable of supporting coldwater fishes*<sup>26</sup>.

Surveys have shown that all three stream types are within the catchment area.

With the size of the area, it is most likely that all the streams have not been found. The following is a listing of the permanent streams that have been located to date. They are named for a connection to a lake, wetland, road or other recognizable location and categorized as cold or warm water. The number of the stream indicates its location on the streams map on the following page.

NAME	Map No.	Classification
Dudman Creek	1	
Wenona West Creek	2	1
Wenona East Creek	3	The initial summer
West Negaunee Creek	4	temperature
Lower Negaunee Creek	5	readings for stream
Upper Negaunee Creek	6	1005.1 - 11 are all 15C or below and
Causeway Creek	7	therefore are
Lost Lake Creek	8	notentially cold
Miskwabi North Creek	9	water streams
Miskwabi Rock Cut	10	Temperature testin
Creek	10	will be continued
Miskwabi Downdraft	11	and recorded.
Creek		
Lily Lake Creek *	12	Warm Water
Burnt River	13	Possible cold water Temperature drops substantially between Long and Loon Lakes.

Figure 13 List of Miskwabi Area Permanent Streams

<sup>&</sup>lt;sup>26</sup> Aquatic Monitoring Program, pg.8



#### Map 13 Miskwabi Area Streams

Credit: MNR "Make a Map" Website

ISSUE: Wetlands and streams in the catchment area of a lake are considered critical to the lake's water quality and general health. ACTION:

- Work to have the five wetlands that scored over 80% on the Haliburton Natural Heritage Mapping Report assessed for Provincial Significance.
- Recommend that waterways (streams and rivers) be added to the County listing of Environmental Protection Zones and included in the Municipal By-laws with the same protection that governs wetlands.
- Work with the Municipality & County to add to the Official Plans 6 wetlands that do not now appear on the Environmental Protection Zones map.
- Recommend that the current Municipal by-law governing development adjacent to wetlands be updated to disallow site alterations to wetlands and streams.
- Recommend that the Municipality support the use of the MNR Environmental Guidelines for Access Roads and Waterway Crossings.

## **FISH COMMUNITY**

The MNR is responsible for the management of fish stocks and fishing regulations in Ontario, including monitoring fish populations, stocking, surveying, protecting habitat, collecting data and longterm planning. The Miskwabi area lakes are under the auspices of the Bancroft District Office. Local field office staff in Minden were very generous with their time and provided our committee access to their testing, stocking and survey information for the four lakes. According to MNR original surveys, a wide variety of minnow species as well as perch and white suckers were found in all four lakes. Also, Wenona, when first assessed in 1958, had a population of Smallmouth Bass. Negaunee was never stocked, but beginning in the 1930's Long, Miskwabi and Wenona were, over the years, individually stocked with one or more of the following species:



*Lake Trout* are freshwater char living mainly in northern North American lakes. About 25% of the world's lake trout lakes are found in Ontario (many in Haliburton County). However, only 1% of Ontario's lakes contain lake trout. Lake trout require deep, cold lakes with well-oxygenated water. They spawn in the fall, at night, scattering their eggs over rocky shoals. The eggs remain among the rocks and hatch in the late winter.

#### Haliburton Gold Lake

*Trout* is a strain of lake trout unique to the Highlands. It has been studied by Dr. John Casselman, a former senior scientist with MNR. His findings indicate that it is one of the oldest and purest strains of lake trout in the world. It appears to be more adapted to



smaller lakes and matures at a young age.

Although lake trout have been a very successful species by adapting to the conditions caused by the last glaciers, they are now potentially vulnerable to conditions of climate changes, invasive species, water level changes and development. This makes the maintenance of the lakes that now have the conditions to support self-sustaining lake trout populations that much more important.

The *Rainbow Trout* is a member of the salmon family and native to the rivers in North America and Asia which flow into the Pacific Ocean. On Canada's west coast, it is called Steelhead. Like salmon, the rainbow is anadromous, meaning it



returns to its original hatching stream to spawn. However, unlike the Pacific coast salmon that die after spawning once, the rainbow can spawn several times. First introduced in 1894 in Ontario, it has become an important sport fish of the Great Lakes and their tributaries, as well as many inland lakes and rivers.



The Smallmouth Black Bass, a member of the Sunfish family, is native to eastern and central North America. It is usually found in moderately shallow water near rocks or submerged logs. They spawn in late spring or early summer when the

male builds a circular nest in gravel. After spawning the male fans the eggs, protects the nest, and guards the young when they are born. This fish is considered by many to be the best-fighting freshwater sports fish.



The Largemouth Black Bass, another member of the Sunfish family, mainly populates the southern part of central North America. Its habitat is the upper levels of the warm water of shallow bays with abundant aquatic vegetation. Nesting is in the early summer and the male guards the nest and young up to one

month after birth. Largemouth Black Bass can grow large – the unofficial record is over 25 lbs. This fish can be distinguished from the Smallmouth Black Bass as its upper jaw extends past its eye.

#### Long Lake

Lake Trout have been stocked in Long Lake on three occasions: 1987, 1991 and 1993.

When last assessed in 1994 MNR stated that the lake only had 13% optimal lake trout habitat and did not warrant further stocking. On the other hand, the assessment did show a large proportion of the population (39%) were natural trout and concluded that both stocked and natural lake trout were moving between Miskwabi and Long Lakes.

As indicated in the Water Quality section, there is reason to be concerned about habitat deterioration for lake trout in Long Lake. Although Total Phosphorus readings are good, Dissolved Oxygen below 10 metres of depth has been on a downward curve. Currently, Long lake is considered "at capacity" for development under the MOE Lakeshore Capacity Assessment, the MNR Dissolved Oxygen criteria and in the Municipality of Dysart et al Official Plan. In 1968, Largemouth Black Bass (199 adults) were stocked in Long Lake (the first and last time) and their progeny are still hitting lures. Smallmouth Black Bass were never directly stocked in Long Lake but migrated from Miskwabi Lake and over the years, especially in the 1970's and 80's, produced a great fishery.

During the 1980's Rock Bass, an invasive species, were unfortunately introduced into Long and Miskwabi Lakes and are abundant today.

As well, over the last five years Pike have been caught in both Long and Miskwabi Lakes.

#### Miskwabi Lake

Miskwabi was first stocked in 1934 with 10,000 lake trout fry. It is thought that these fish did not prosper and between 1965 and 1977 20,000 lake trout were introduced. From all accounts these lake trout produced a fishery that was second to none in the Highlands during the 1970's and 1980's. The following is a chronological listing of some interesting details from the MNR records regarding Miskwabi Lake:

- 1970 Department of Lands and Forests fact sheet indicates a strong lake trout fishery.
- 1980 MNR assessment states Miskwabi is the prime lake trout lake in Minden District. "The best producer of lake trout in terms of pounds per acre ...produces at least 3.9 times more lake trout per hectare of water than any other lake surveyed in the Minden District."<sup>27</sup>
- 1986 Assessment of habitat conditions, spawning beds and stocked vs. naturally reproducing fish. Results indicated that the natural fish made up 87% of the population. Miskwabi was estimated as having 61% optimal habitat (volume of the lake colder than 10°C and >6 mg/L of Dissolved Oxygen). The report recommended that Miskwabi be managed as a naturally selfsustaining lake trout lake.
- 1988 Assessed with recommendations for improving natural lake trout populations.
- 2000 First stocking of Haliburton Gold lake trout in Miskwabi.
- 2000-2010 Dr. David O. Evans, research scientist of MNR's Aquatic Research and Development Section, initiates Haliburton Lake Trout Adoptive Management Study (2000 – 2010). Miskwabi was selected as one of the four Haliburton lakes used to evaluate the effects of slot-size regulations and stocking using Haliburton Gold on fishing effort, catch, harvest and population responses of the exploited lake trout stocks. For Miskwabi, Haliburton Gold was stocked and the slot size for lake trout was removed.

Although Dr. Evans' results have not yet been formally published, the MNR website summarizes the findings of the study:

- "Slot-size regulations By reducing fishing mortality, researchers saw that slot size regulations clearly protected lake trout of prime spawning ages (6 to 10 years). This shows that regulations are an effective management approach for ensuring the long-term sustainability of wild lake trout stocks.
- No slot-size regulations After slot size regulations were removed in two lakes (including Miskwabi), the rate of people catching and keeping their fish dramatically increased during the first few years. The total harvest of fish then fell back to pre-slot amounts showing that high fishing pressure has depleted these populations.
- Stocked and no size limits Stocking in the absence of a slot-size regulation provided some incentive for anglers but could not counteract the effects of high fishing pressure and high mortality rates, placing the population at risk.
- Stocked...but with size restrictions Stocking has not proved to be a "hook" for anglers when slot-size regulations are in place, because what anglers are allowed to keep (based on size) is restricted to provide protection for the highly vulnerable, mature adults."<sup>28</sup>

The report concludes that "a management strategy that combines slot-size protection with stocking of lakes with poorer habitat seems to be the best approach for providing both angling opportunities and ensuring the long-term future for Ontario's native lake trout."<sup>29</sup> With regard to habitat and water quality, Miskwabi Lake is well suited for natural trout populations. It is buffered from acid rain and has strong calcium levels which allow for diverse and abundant phyto- and zooplankton populations necessary for the survival of young lake trout. As well, the Dissolved Oxygen readings continue to be very good although the bottom readings (40m) are declining.

<sup>&</sup>lt;sup>27</sup> Miskwabi Lake Fisheries Report 1980, pg.2

 <sup>&</sup>lt;sup>28</sup> www.mnr.gov.on.ca/en/Business/Aquatics. June 16, 2013
<sup>29</sup> ihid

Under the MOE Lakeshore Capacity Assessment the lake is at capacity and this is reflected in the Dysart et al Official Plan. For this reason, when the north shore of the lake was developed in the 1990's the number of building lots was reduced by about 60%, lot sizes were substantially increased, building setbacks were increased and special rules introduced regarding tree removal. As well, four environmental zones were created to protect fish habitat.

#### Negaunee Lake

According to MNR files Negaunee has never been stocked with fish. It was first assessed by the Department of Lands and Forests in 1958 with the notation that there were an "exceptionally large number of minnows of all sizes up to 6 inches".<sup>30</sup> As well, the lake was measured to be 23.75 m (78 ft.) deep with good temperature gradient. At the time, the District Biologist indicated that "the lake should be particularly suitable for speckled trout". A creel census in 1984 indicated a large number of perch and minnows being caught, but no game fish. Today, residents of the lake indicate that there is a self-sustaining Smallmouth Bass population in the lake.

#### Wenona Lake

When first assessed in 1958, Wenona was found to have a heavy population of Smallmouth Bass and a variety of non-game species including perch, pumpkinseed, suckers and minnows.

Based on an assessment in 1971 that found Wenona to be 19.2 m (63 ft) deep with good temperature stratification and reasonable dissolved oxygen, lake trout were stocked on four occasions between 1972 and 1979. Unfortunately, that did not produce a sustainable fishery.

In 1988 Largemouth Bass were stocked in the lake and have continued to survive to this day.

Starting in 1999 Rainbow Trout were introduced to Wenona and

between that time and the present, 20,000 fish have been stocked. This has produced fishing excitement for many over the years. As there are no streams entering Wenona suitable for spawning, the rainbows do not reproduce.



Ice Fishing on Miskwabi Photo: Ken Morrison

ISSUE: It is well recognized that lake trout populations are in peril in Haliburton. ACTION:

- Support the recommendations of Dr. David Evans regarding the importance of the release of trout caught within restricted size (slot sizes) and the protection of lake trout lakes (like Miskwabi Lake) that have good habitat for natural lake trout reproduction.
- Support the CHA initiative, now with the MNR, to close the lake trout season earlier to protect breeding trout.
- Encourage a "catch and release" policy for lake trout on Miskwabi and Long lakes.
- Encourage the use of lead-free fishing equipment.
- Work with MNR to rejuvenate self-sustaining lake trout populations in Long and Miskwabi Lakes.

<sup>&</sup>lt;sup>30</sup> Department of Lands & Forests memorandum 1958

## WILDLIFE AND WILDLIFE HABITAT

The Miskwabi area is surrounded by over 3000 square kilometres of relatively pristine wildlife habitat. It includes many lakes, rivers, large wetlands and abundant highlands. The landscape is bordered on the south by Hwy. 118, the west by Hwy 35, the east by Hwys. 62/127 and the north by Hwy. 60, but not intersected by any. This gives wildlife an extensive, natural ecosystem with good travel corridors extending into Algonquin Park.

#### Mammals

The chart on the right lists the mammals of the Haliburton Highlands. The Miskwabi area has a population of most, if not all. Some mammals of note include:



The *Eastern Wolf* is now recognized as a distinct subspecies of the Grey Wolf. Prior to European contact the Eastern Wolf and the Red Wolf of the Eastern U.S. were likely the same species. Owing to habitat fragmentation, hybridization with the Coyote and Grey Wolf, and

Eastern Wolf on Miskwabi Photo: William Bunn

genetic drift, the Eastern Wolf and Red Wolf now exhibit genetic differences. There are probably fewer than 500 Eastern Wolves in Canada with the highest population found in Algonquin Park. Wolf packs require relatively large, unbroken forests with home ranges as large as 500 sq. kms. The Eastern Wolf is listed as a species of Special Concern under Ontario's Endangered Species Act.<sup>31</sup>

Haliburton Highlands Mammals <sup>32</sup>			
Beaver	Northern Long-eared Bat		
Big Brown Bat	Northern Short-tailed Shrew		
Black Bear	Muskrat		
Coyote	Porcupine		
Deer Mouse	Pygmy Shrew		
Eastern Chipmunk	Raccoon		
Eastern Cottontail	Red Fox		
Eastern Pipistrelle	Red Squirrel		
Eastern Wolf	River Otter		
Ermine	Rock Vole		
Fisher	Silver-haired Bat		
Grey Squirrel	Small-footed Bat		
Hairy-tailed Mole	Smoky Shrew		
Hoary Bat	Snowshoe Hare		
Least Chipmunk	Southern Bog Lemming		
Least Weasel	Southern Flying Squirrel		
Little Brown Bat	Southern Red-backed Vole		
Long-tailed Weasel	Star-nosed Mole		
Lynx	Striped Skunk		
Marten	Virginia Opossum		
Masked Shrew	Wapiti – (Elk)		
Meadow Jumping Mouse	Water Shrew		
Meadow Vole	White-footed Mouse		
Mink	White-tailed Deer		
Moose	Woodchuck		
Northern Flying Squirrel	Woodland Jumping Mouse		

Figure 14 List of Mammals

<sup>&</sup>lt;sup>31</sup>Species at Risk. www.mnr.gov.on.ca

<sup>&</sup>lt;sup>32</sup> County of Haliburton Natural Heritage Mapping, pg.53

Another species of note is the *Moose*. According to the Ministry of Natural Resources, the Haliburton Moose population has been increasing since 1998 and is now within the desired ecological density of 20 - 40 moose per 100 km<sup>2</sup>.



Photo: Terry Anderson

Moose travel over large areas and have seasonal ranges. The chart below shows their seasonal preferred habitat.



Figure 15 Seasonal Movements of Moose in Ontario Credit: Moose Resource Report

The large green zone around Miskwabi Lake and to the north-east in the map below indicates a Moose late Wintering Area as designated by MNR.



Map 14 Moose Late Wintering Area

Credit: County of Haliburton

The expanse to the north and west sides of Miskwabi Lake is considered preferred habitat for the *Southern Flying Squirrel*.

Also of interest is the *Cougar* which by the early 20<sup>th</sup> Century was mainly confined to the Rockies and the West Coast. Due to conservation efforts during the past 40 years the Cougar has moved eastward. Are they extending their range into central and southern Ontario in pockets of undisturbed forest with healthy deer populations? Over the last five years, on different occasions, four members of our community have reported seeing a cougar.

### Birds

The Miskwabi area, with its mixed habitat of hardwood uplands, coniferous lowlands, lakes and wetlands, is ideal for a large variety of bird species. Many migrate long distances from Central and South America; whereas, others are year-round residents.

The following chart outlines a number of uncommon birds that inhabit the spaces around the lakes. Reference is given to the preferred habitat of each and the likelihood of each species breeding in this region.

Common Name	Natural Heritage Mapping – Preferred Habitat	Breeding Bird Atlas of Ontario Indications of Breeding <sup>33</sup>
Red-shouldered Hawk	Deciduous forest	Confirmed
Veery	Deciduous forest	Probable
Wood Thrush	Deciduous forest	Probable
Canada Warbler	Deciduous forest	Confirmed
Black-throated Blue Warbler	Deciduous forest	Probable
Northern Cardinal	Deciduous forest	Probable
Blackburian Warbler	Coniferous forest	Probable
Black-throated Green Warbler	Coniferous forest	Confirmed

Figure 16 Uncommon Birds of Miskwabi Area

One particularly rare species is the *Golden-winged Warbler*. This beautiful song bird nests in eastern-central North America and migrates to Central and South America in the winter. Its numbers are in decline (it is believed caused by habitat loss) and it is listed as a species of Special Concern in the Ontario Endangered Species Act. The Breeding Bird Atlas of Ontario indicates that according to their research, the Golden-winged Warbler does not nest in our area. The picture of a Golden-winged Warbler below, taken on Miskwabi Lake in May 2014, might be of one migrating through the area, or possible evidence that there are some breeding pairs around the Burnt River headwaters.



Golden-winged Warbler Photo: William Bunn

As well, Larry Holden, a long-time resident, has confirmed sightings of the young of the *Canada Goose, American Woodcock, Evening Grosbeak and Wild Turkey* even though these birds are not confirmed as breeding in the area according to the Atlas of the Breeding Birds of Ontario Chart on pg.48.

Also of note, in recent years, *Bald Eagles* have spent the winter in the area of the lakes and there have been reported sightings of *Golden Eagles*. The *Double-crested Cormorant*, as it extends its range, has made appearances on the lakes in the past few summers which is a concern as their colonies can kill trees and vegetation.

<sup>&</sup>lt;sup>33</sup> Breeding Bird Atlas of Ontario website

The *Common Loon* is one of the birds most associated with cottaging. They are seen and heard on all four lakes and normally have successful young on Long, Miskwabi and Wenona Lakes. In the case of Wenona and Miskwabi, the nests have been built on platforms installed and maintained by local property owners. Ideally, loons nest on peninsulas and islands as the set-back placement of their feet makes walking on land difficult. Platforms provide easy access which might be otherwise threatened by lowered water levels occurring during the early-June to mid-July nesting season. On Miskwabi and Wenona Lakes platforms have helped produce very successful breeding rates over the years. According to Bird Studies Canada<sup>34</sup>, an average reproductive success of one young every other year is considered good. Based on this, Miskwabi, Long and Wenona Lakes would be considered very successful loon lakes. Miskwabi, surveyed for the Canadian Lakes Loon Survey since 1995, has reported better than average results. Frequent sightings of large young on Long and Wenona Lakes late in the season would also confirm their reproductive success rates.



Day old baby loon July 2014 on Miskwabi Lake platform

Photo: Mike Tipping



Loon family on Miskwabi July 2013 Photo: Mike Tipping

Bird Studies Canada reports the reproductive success of loons is still declining in Ontario due to lingering effects of mercury and acid rain precipitation. As the geology of our area has protected the lakes from the effects of acid rain, this could be a second reason for the reproductive success.

The Atlas of the Breeding Birds of Ontario (ABBO) is an outstanding document of bird research in Ontario. It is based on 150,000 hours of field time and 1.2 million individual breeding bird records. The atlas divides the province into 10km<sup>2</sup> areas and details the current distribution of each breeding species in the province and how its distribution has changed over the past 20 years. The table on the following page lists the results for the Miskwabi area.

<sup>&</sup>lt;sup>34</sup> Canadian Lakes Loon Survey 1981-2012, pg.5

Figure 17 Atlas of the			
Breeding Birds of Ontario			
Broading Fridance			
in Mielewshi Are			
Alder Elvesteher	Probablo		
American Bittorn	Possible		
American Block Duck	Possible		
American Crow	Confirmed		
American Coldfineh	Probablo		
American Goldmich	Probable		
American Redstart	Confirmed		
American Woodcool	Prohable		
Relting and Origin	Descible		
Baltimore Uriole	Possible		
Bank Swallow	Confirmed		
Barn Swallow	Confirmed		
Barred Owl	Probable		
Belted Kingfisher	Confirmed		
Black-and-white Warbler	Confirmed		
Black-backed Woodpecker	Possible		
Black-billed Cuckoo	Possible		
Blackburnian Warbler	Probable		
Black-capped Chickadee	Confirmed		
Black-throated Blue Warbler	Confirmed		
Black-throated Green Warbler	Confirmed		
Blue Jay	Confirmed		
Blue-headed Vireo	Probable		
Broad-winged Hawk	Confirmed		
Brown Creeper	Confirmed		
Brown-headed Cowbird	Possible		
Canada Goose	Possible		
Canada Warbler	Confirmed		
Cedar Waxwing	Probable		
Chestnut-sided Warbler	Confirmed		
Chipping Sparrow	Confirmed		
Common Grackle	Confirmed		
Common Loon	Confirmed		
Common Merganser	Confirmed		
Common Nighthawk	Possible		
Common Raven	Possible		
Common Snipe	Probable		

Common Yellowthroat	Confirmed
Downy Woodpecker	Possible
Eastern Bluebird	Confirmed
Eastern Kingbird	Probable
Eastern Phoebe	Possible
Eastern Towhee	Possible
Eastern Wood-Pewee	Confirmed
European Starling	Confirmed
Evening Grosbeak	Probable
Golden-crowned Kinglet	Probable
Gray Catbird	Possible
Gray Jay	Possible
Great Blue Heron	Confirmed
Great Crested Flycatcher	Probable
Great Horned Owl	Probable
Green-winged Teal	Probable
Hairy Woodpecker	Confirmed
Hermit Thrush	Probable
Herring Gull	Possible
Hooded Merganser	Probable
House Sparrow	Possible
House Wren	Possible
Indigo Bunting	Confirmed
Killdeer	Possible
Least Flycatcher	Probable
Lincoln's Sparrow	Possible
Magnolia Warbler	Confirmed
Mallard	Probable
Marsh Wren	Possible
Merlin	Possible
Mourning Dove	Probable
Mourning Warbler	Probable
Nashville Warbler	Confirmed
Northern Cardinal	Probable
Northern Flicker	Possible
Northern Goshawk	Possible
Northern Parula	Possible
Northern Saw-whet Owl	Possible
Northern Waterthrush	Probable
Olive-sided Flycatcher	Possible
Osprey	Confirmed

Ovenbird	Confirmed
Philadelphia Vireo	Possible
Pileated Woodpecker	Probable
Pine Warbler	Probable
Purple Finch	Probable
Red-breasted Nuthatch	Confirmed
Red-eyed Vireo	Confirmed
Red-headed Woodpecker	Possible
Red-shouldered Hawk	Probable
Red-tailed Hawk	Possible
Red-winged Blackbird	Probable
Ring-necked Duck	Probable
Rock Pigeon	Confirmed
Rose-breasted Grosbeak	Confirmed
Ruby-crowned Kinglet	Possible
Ruby-throated Hummingbird	Confirmed
Ruffed Grouse	Confirmed
Savannah Sparrow	Probable
Scarlet Tanager	Probable
Sharp-shinned Hawk	Possible
Song Sparrow	Probable
Spotted Sandpiper	Confirmed
Swainson's Thrush	Possible
Swamp Sparrow	Probable
Tree Swallow	Confirmed
Turkey Vulture	Possible
Veery	Probable
Warbling Vireo	Confirmed
Whip-poor-will	Possible
White-breasted Nuthatch	Confirmed
White-throated Sparrow	Confirmed
White-winged Crossbill	Confirmed
Wild Turkey	Possible
Willow Flycatcher	Possible
Winter Wren	Confirmed
Wood Duck	Probable
Wood Thrush	Probable
Yellow Warbler	Probable
Yellow-bellied Sapsucker	Confirmed
Yellow-rumped Warbler	Confirmed
Yellow-throated Vireo	Probable

### **Reptiles and Amphibians**

The numerous wetlands, riparian zones and shorelines in the Miskwabi Lakes area offer habitat for a variety of reptiles and amphibians. In Ontario, the populations of many reptiles and amphibians have declined significantly over the past 30 years. The reasons are not fully understood, but habitat loss is considered to be a major factor in the decline.

The following is a list of herpetofaunal (reptiles and amphibians) whose range includes our area of Haliburton County:

Haliburton Herpetofaunal <sup>35</sup>			
Turtles	Amphibians		
Blanding's Turtle	Frogs and Toads		
Midland Painted Turtle	American Toad		
Snapping Turtle	Bull Frog		
Spotted Turtle	Green Frog		
Stinkpot Turtle	Grey Treefrog		
Wood Turtle	Leopard Frog		
Snakes	Mink Frog		
Dekay's Brownsnake	Pickerel Frog		
Eastern Gartersnake	Spring Peeper		
Eastern Hog-nosed Snake	Western Chorus Frog		
Eastern Milksnake	Wood Frog		
Eastern Ribbonsnake	Salamanders and Newts		
Eastern Smooth Greensnake	Blue-spotted Salamander		
Northern Red-bellied Snake	Eastern Red-backed Salamander		
Northern Watersnake	Four-toed Salamander		
Ring-necked Snake	Northern Two-lined Salamander		
	Red-spotted Newt		
	Spotted Salamander		

Figure 18 Haliburton Herpetofaunal



Blanding's Turtle





Blue-spotted Salamander

**Spring Peeper** 



Northern Watersnake

<sup>&</sup>lt;sup>35</sup> County of Haliburton Natural Heritage Mapping, pg.51

### **SPECIES AT RISK**



Snapping Turtle born at Miskwabi Lake Causeway Sept. 2013 Photo: Michael Giza

The picture above of a *Snapping Turtle* that has just hatched in the gravel at the side of a Miskwabi Lake road depicts a species at risk. Habitat depletion is one of the main threats but other important causes, in particular for reptiles, include road kill and illegal harvesting. This makes educational efforts and habitat stewardship essential in helping to prevent the decline of these species. The Species at Risk in Ontario (SARO) listing is produced by the Ontario Ministry of Natural Resources to give status designations for species at risk in the province.

The table opposite is a list of a number of species whose range includes the Miskwabi Lakes area and which are on the SARO (updated May 2013) list.

Miskwabi Lake	es Area Species at Risk
Endangered *	Cougar
	Golden Eagle
	Little Brown Rat
	Wood Turtle
Threatened **	Blanding's Turtle
	Eastern Hog-nosed Snake
	Spotted Turtle
	Stinkpot Turtle
	Whip-poor-will
Special Concern ***	Bald Eagle
	Canada Warbler
	Common Nighthawk
	Eastern Milksnake
	Eastern Ribbonsnake
	Eastern Wolf
	Monarch Butterfly
	Snapping Turtle

Figure 19 Miskwabi Area Species at Risk

\*Endangered – lives in the wild but is facing imminent extinction or extirpation.

\*\*Threatened (THR) - lives in the wild, is not endangered, but is likely to become endangered if steps are not taken to address factors threatening it.

\*\*\*Special Concern (SC) – lives in the wild, is not endangered, but may become threatened or endangered if steps are not taken to address factors threatening it.

# **INVASIVE SPECIES**



Invasive species can be those that are exotic (non-native), and those that are native to Ontario but not native to the local area. The introduction of such species to a local ecosystem can bring widespread, unpredictable, and in many cases detrimental effects to habitats. Being at the

headwaters of the Burnt River system, the lakes cannot be easily affected by upstream invaders. On the other hand, we have the additional responsibility of not introducing invaders to the headwaters of the system.

Invasive species that have become a concern in the Haliburton Highlands can be categorized as follows:

 Aquatic animals – Zebra Mussels, Spiny Water Flea (pictured above) Rusty Cray Fish, Rock Bass, and Round Goby have all been introduced to many water bodies. All the Miskwabi area lakes have been tested for years through involvement in the Invading Species Watch Program of the MNR and the OFAH. As of the 2013 testing, the four lakes are free of Zebra Mussels and Spiny Water Fleas.

As boats and bait buckets are the main means of transferring invasive species, it is essential that the steps listed below are followed to keep the lakes free of these species.

- 1. Drain all water from bilges and wells on land.
- 2. Empty bait buckets on land.
- 3. Never release live bait.
- 4. Inspect, wash and dry boats, trailers and equipment before launching in a new water body.
- Aquatic Plants Purple Loosestrife, European Frogbit, Eurasian Milfoil, Fanwort and Flowering Rush are affecting Haliburton lakes. To date, we have not confirmed any of these invaders.

 Cottage plants – Cottage country is susceptible to the introduction of plants such as Goutweed and Periwinkle that can be brought from suburban areas. Here they compete with native species, including tree seedlings, for light, nutrients and water. Goutweed is highly shade-tolerant and competitive once established, reproducing by seed and by spreading underground. Its vigorous nature can choke out native groundcovers. Periwinkle also thrives in dry shade and has few pests or diseases outside its native range of Europe.

The Ontario Invasive Plant Council created the "*Grow Me Instead Campaign*" to inform gardening enthusiasts about popular invasive plants that can cause problems in the landscape. They highlight a variety of non-native plant alternatives found to be non-invasive.<sup>36</sup>

#### **Invasive Species Mapping**

Early Detection & Distribution Mapping System (EDDMapS) is a web-based mapping system launched in 2005 by the University of Georgia to document invasive species distribution. The Ontario programme was developed through support and funding provided by Canadian/Ontario Invasive Species Centres, the OFAH and the MNR. Numerous agencies and monitoring programmes are in place to collect information. Programme involvement is now being offered to those now participating in the Invading Species Watch.

#### ISSUE:

Invasive species can cause severe, negative alterations to natural ecosystems as well as be troublesome to people. ACTION:

- Continue to work with the *Invading Species Watch* programme.
- Increase educational material and signage to prevent contamination of the lakes by boats and bait buckets.
- Have MACA join the new MNR/Federation of Ontario Cottagers Association (FOCA) EDDMapS programme.

<sup>&</sup>lt;sup>36</sup> More at <u>www.ontarioinvasiveplants.ca</u>.

# PHYSICAL ELEMENTS

Physiography refers to the physical geography of an area involving the landscape, its various landforms, makeup and gradients. The Miskwabi area is located in the Pre-Cambrian Shield physiographic region which is characterized by granite bedrock with thin overburden. The area's bedrock is mildly fractured, moderately dipping granite gneiss and often found as steep cliffs or outcroppings along the shorelines.

# SOILS

The soil substrates are generally thin and shallow with some areas of heavier glacial gravel deposits. The topography of the drainage basin is steep to gently rolling hills which have a maximum elevation of approximately 80 m above the level of the lakes. Unusual for the Haliburton Highlands, the lakes of the Miskwabi area have good calcium levels due to the composition of the surrounding geology. For this reason the lakes are not threatened by acid rain and have strong zooplankton populations.

# MINERALS AND AGGREGATES

There are no mining claims or aggregate operations in the area of our lakes. It is well understood that mining and aggregate operations in proximity to wetlands, streams or lakes can have a detrimental effect on the water quality and natural environment.

# **STEEP SLOPES**

Developments in areas of steep slopes can result in unwanted alterations to the natural landscape, negative visual impact, as well as increased erosion, slope instability, heavier storm water run-off and potential damage to fish and wildlife habitat. Steep slopes are present on all four lakes, but particularly on Long and Miskwabi Lakes. During the years of development these presented a substantial challenge in completing the required infrastructure while at the same time maintaining natural and environmentally-friendly landscapes.

# NARROW WATER BODIES

There are three narrow waterways in the lakes, all with lakefront lots. The first is located at the narrows between Long and Miskwabi Lakes and the other two are farther to the west on Long Lake. They require special caution when used by motorized boats for three reasons: safety concerns; helping to reduce shoreline erosion; and minimizing disturbance to the lakefront owners. Stewardship efforts are required to encourage all motorized boat operators using these narrow zones to maintain the legal 10km/hr (6mph) or less boat speed within 30 metres (100ft) of shorelines.



Long Lake Narrows Photo: Andy Mosher

# SOCIAL ELEMENTS

# SOCIAL CULTURE

According to the Merriam-Webster dictionary, the definition of the word "social", is "relating to or involving activities in which people spend time talking to each other or doing enjoyable things with each other". For the purpose of our watershed plan we could also say there is a "social" involvement with people and the environment.

Our four lakes do not have the reputation of being "party" communities like some others in the area. People prefer to "do their own thing" with family, friends and lake neighbours. However, most property owners are very community conscious which is attested to by the active involvement by residents and cottagers over the past 40 years in the Miskwabi Area Community Association, formerly the Miskwabi Area Cottagers' Association.



MACA 40<sup>th</sup> Anniversary AGM July 5, 2014

Photo: Susanne James

Over half of the property owners on the lakes, back lots and roadways are currently members of the association, some of them since 1974.

The association has an environmental as well as social agenda: planned activities range from an annual road clean-up and BBQ held in May to an annual association meeting and BBQ held in July to give members the opportunity to share issues and concerns for the directors to consider and address on their behalf. As a result of this type of input, and reflecting the stakeholder survey results, MACA has developed a new "Owners, Renters and Visitors Guide" to remind all of the respect needed for the social and environmental community.

# LANDSCAPE AND AESTHETICS

Survey respondents clearly indicated the importance of maintaining the special nature of the four lakes. In the Natural Features section, "Lakes" scored 100% as Very Important/Important. "Forests and Shorelines" came a close second and third. Development can seriously compromise the enjoyment of these values. The water's edge is impacted by the building of waterfront structures or the removal of vegetation. The skyline and is artificially altered by the construction of buildings, power lines or communication towers.

# TRANQUILITY AND NIGHT SKIES

A natural dark sky was rated as Very Important/Important by over 80% of respondents as adding to their enjoyment of the lakes. Outdoor lighting adds to the enjoyment of evening activities and provides safety. Both needs can be met if lighting is kept to a minimum, installed to light only local areas, and off when not in use. Over 80% of respondent cited "Peace and Quiet" as important to their enjoyment of cottage life. Fireworks have been cited in a MACA survey as being a source of light and noise pollution and the majority of respondents to that survey thought they should be limited to statutory holiday events. Again, respect and consideration for others is important. Safety and the possible starting of forest fires, as well as the prospect of chemicals and debris dropping in the water are issues of concern, as are the noise effect on humans, wildlife and the environment. While survey results were made available to the Municipality, at this time there are no municipal by-laws regulating the sale or use of fireworks. Dysart et al does have a noise pollution by-law that requires noise levels to be reduced after 11p.m.; however, enforcement is difficult.

#### **ISSUE:**

Over 80% of survey respondents indicated that "peace and quiet" and "night skies" were "Very Important" or "Important" to them.

### ACTION:

- Support the educational handout *Guidelines for Owners, Renter and Visitors* developed by MACA. The one-page document is available for all to use.
- Develop separate educational handouts for owners who rent and renters. These would cover the environment, septic systems, waste disposal, noise and other topics related to renting.

# **RECREATIONAL ACTIVITIES**

People on the lakes enjoy a wide range of activities – land or water based.

Most important seemed to be non-motorized ones: land-based (walking, hiking, bird watching) or water-based (swimming, canoeing/kayaking/sailing or fishing) as evidenced by preferences cited in the stakeholder's survey.

#### Land-based

From the survey, snowmobiling is an activity "Very Important/Important" to approximately 40% of respondents. Haliburton County has an active snowmobile association (HCSA). It has a large involvement in trail grooming and 370 km of groomed trails with spectacular scenery. There are also trails maintained in close proximity to the four lakes that are groomed by the Paudash Trailblazers Snowmobile Club (east and north behind Miskwabi Lake, and southeast of Negaunee Lake). There are no trails marked over ice-covered lakes and few alongside the roadways included in this watershed plan.

ATVing is also "Very Important/Important" to approximately 30% of respondents. Currently ATVers also use snowmobile trails.

#### Water-based

Swimming is the most important water activity; enjoyed by 98% of respondents. Another activity receiving a high rating of importance was fishing. As set out in the recommendations in the Natural Features section, encouraging a catch and release policy and the use of lead-free equipment could help ensure an active fishery for future generations.



Long Lake Catch Photo: Phil Wiseman

# **RECREATIONAL BOATING and SAFETY**

Residents and visitors to the four lakes enjoy many types of activities in watercraft; from kayaking, to fishing in a "tinny", to skiing behind a wakeboard boat.

Property owners and visitors can make use of two public launch points: one at the end of Wenona Lake and the other at the "airstrip" on Miskwabi Lake. Of note though, through common agreement motorized boats are not used on Negaunee due to its small size.



Tubing on Miskwabi Lake

Photo: Virginia Vranckx

Like most lakes, the Miskwabi area lakes have seen an increase in the number and type of boats over the years. Newer boats are often pontoon type used to tour the lakes or pull tubes. There are several wakeboard boats and personal watercraft. The latter, as well as any boat driven at high speed too close to shore, have recently caused concern to property owners along narrow areas of Long Lake and the channel between Long Lake and Miskwabi. The effects of large wakes and inappropriate boat speeds can damage the environment and cause safety concerns. Large wakes travel long distances and increase the potential for damage to floating docks and possible injury to people on them. As well, the erosion of shoreline affects spawning beds and weed beds resulting in the loss of fish habitat and encouraging the spread of invasive species.

Responsibility rests with residents and visitors to report serious infractions to the Ontario Provincial Police (OPP) Marine unit as the OPP's presence is restricted to occasional visits to the lakes. Currently signs regarding boat speed near shoreline are posted only at the boat launches and the narrows between Miskwabi and Long Lakes.

# RECREATIONAL BOATING CARRYING CAPACITY

Recreational carrying capacity was defined in the 1970's as:

"the ability of a water body to accommodate particular types of uses and a given number of users within defined levels of ecological disturbance and inter use conflicts."<sup>37</sup>

It refers to the number of watercraft that can simultaneously operate on a lake without:

- compromising user safety;
- causing significant user displacement and dissatisfaction;
- causing environmental harm to the resource;
- and/or, causing significant dissatisfaction to non-boaters.

<sup>&</sup>lt;sup>37</sup> Kusler, pg.7

This topic has not yet received much study in Canada but has been analyzed by a number of studies in the United States. We believe the subject is an important one and should not be left unreviewed.

For Haliburton County, recreational carrying capacity is relevant for at least three reasons:

- 1. The Highlands are within easy access of the burgeoning metropolitan populations of southern Ontario.
- 2. Lakes in Ontario are a public resource without restriction to the number and the size of boats.
- 3. In general, the lakes of Haliburton County do not have large surface areas. Smaller lakes usually have relatively less surface area in relation to the perimeter than do larger lakes.

The following table demonstrates this using a perimeter factor of 33 metres representing an average lakefront lot.

COMPARISON OF LAKE SURFACE AREA and PERIMETER					
LAKE	SURFACE AREA (acres)	PERIMETER (metres)	LAKE SURFACE acres per 33m lot		
Long	218	10620	.68		
Wenona	146	5500	.88		
Miskwabi	662	10600	2.1		
Kennisis	3501	41000	2.8		
Muskoka	29990	269000	3.7		
Simcoe	716358	303000	78.0		
Ontario	4,685,114	1,146,000	134.9		
Figure 20 Compa	ricon of Lake Surface Are	a and Darimatar	Courses MAND data		

Source: MNR data

Development around smaller lakes can more rapidly impact safe and enjoyable boating space. This was demonstrated in Wisconsin by the Lake Ripley (418 acres) Watercraft Census and Recreational Carrying Capacity Study of 2003.

#### Calculating Carrying Capacity

- 1. Carrying capacity is determined by dividing the Usable Lake Surface Area by Optimum Boating Density.<sup>38</sup>
- 2. Usable Lake Surface Area is calculated by starting with the lake's surface area and deducting a shoreline buffer, narrow passages, shallow areas, and environmentally sensitive areas.
- 3. Optimum Boating Density is the surface area per watercraft required for boating activities. This can be viewed as the space required for motorized, non-motorized, and all activities combined. As human perception is involved in judging overcrowding, enjoyment and safety levels, there is no universally accepted Optimum Boating Density. Using the figures in nine U.S. reports we calculated that the average minimum space required for combined boating activities is 14.4 acres per boat.<sup>39</sup>
- 4. Calculate the Recreational Carrying Capacity of the lake by dividing the Usable Lake Area by the Optimum Boating Density.
- 5. Compare the Recreational Carrying Capacity to the actual use.40

<sup>&</sup>lt;sup>38</sup> Recreational Carrying Capacity in Lakes: How much is too much?

<sup>&</sup>lt;sup>39</sup> Techniques for Estimating Boating Carrying Capacity, pgs.7, 22.

<sup>&</sup>lt;sup>40</sup> Two actual use surveys were completed for each lake on June 28 & 29, 2014.

The following table, using the above method, gives the results for 3 of the 4 lakes included in this watershed plan.<sup>41</sup> For Usable Surface Area we did not deduct a shoreline buffer, narrow passages, shallow areas, and environmentally sensitive areas which, when calculated, on average totaled 33% of a lake's surface. The reason is, in our survey observations, it was noted that non-motorized boats make use of the total surface area of a lake.

RECREATIONAL BOAT CARRYING CAPACITY TABLE					
LAKE	Total Surface Area (acres)	Optimum Boating Density (acres)	Carrying Capacity	Boat Use Survey Count	+/- Optimum Capacity
LONG	218	14.4	15	26	+58%
MISKWABI	662	14.4	46	33	-39%
LONG - MISKWABI COMBINED	880	14.4	61	59	-3%
WENONA	146	14.4	10	13	+30%

Figure 21 Recreational Boat Carrying Capacity Table

On many summer days all the lakes appear to be quiet; however, as calculated above, during peak use periods both Long and Wenona Lakes exceed their Optimum Boat Carrying Capacity. As well, if the figures for Long and Miskwabi Lakes are combined (as the two lakes are connected), the combined use approaches the Optimum Capacity. This is important for the future as trends indicate:

- a. More boats per property
- b. More large, powerful or fast watercraft
- c. Increased variety of lake activities
- d. Higher percentage of rental cottages
- e. Greater use of public boat launch facility

#### ISSUE:

With swimming enjoyed by 98% of survey respondents, various forms of boating by 80%, faster and more powerful watercraft, and crowded weekend conditions, safety has become a growing concern. This concern is supported by a review we conducted of U.S. studies regarding recreational carrying capacity of lakes.

#### ACTION:

- Develop a Boating Code of Conduct for property owners and lake users explaining the rules for safe and pleasurable operation including following the Safe Boating Guide, minimizing wake near shores and narrows, respect for neighbours, use of clean boats and trailers for environmental reasons, and a no alcohol on-board policy.
- In addition to the slow speed sign at the narrows between Long and Miskwabi Lakes, add a new sign at each of the narrows farther to the west on Long Lake.
- Recommend that the Municipality and County consider using a Recreational Carrying Capacity Model as one of the criteria when considering new or increased development on/adjacent to lakes.

<sup>&</sup>lt;sup>41</sup> Negaunee Lake was excluded as it has no public boat launch and a selfregulated no-motorized boat policy.

# LAND USE

This section of the Watershed Plan provides a description of the current land uses around the lakes as well as the various policies, provisions and zoning by-laws that govern land use.

### **SUMMARY**

Miskwabi area watershed is characteristic of typical cottage country. As such, it is used predominantly for residential and recreational purposes with most of the development occurring along the shoreline of the four main lakes.



Aerial View of Miskwabi Lake North Shore.

Photo: Mike Tipping

In fact, all land in the region is zoned for use as residential, rural or open/environmentally protected space. There is no commercial, industrial or institutional land use in the region.

There are also no large parcels of Crown Land. This is because the Crown sold the entire area to the Municipality in the 1860s.

Forestry has and continues to be an important economic activity and source of employment within the Municipality. There continues to be modest logging and forestry activity occurring in the watershed. Land uses are governed by zoning regulations administered by the Municipality of Dysart et al (as described below).

In the early 1960's there was very little development on the lakes, but growth of seasonal and permanent residences has been steady over the past five decades. As of January 2014, there are 385 occupied waterfront properties and 66 vacant waterfront lots. There has also been a steady growth of backlot residences. Nine separate Environmentally Protected areas are connected to the main four lakes.

There are also a number of important references to confirm that Miskwabi and Long Lakes are at capacity (as defined by the Province of Ontario Lakeshore Capacity Assessment Handbook and as referenced in the Dysart et al Official Plan).

As such, creation of additional building lots and the associated development could seriously jeopardize lake trout populations and water quality.

Based on the Dysart et al Municipal zoning maps and manual measurements, a land use profile for the MACA lakes was created. The data shows that the four main lakes have approximately 29 kms of waterfront. 25% of the waterfront is zoned Open Space or Rural and therefore should see little development. Approximately 70% of properties are served by serviced roads and therefore could more likely become permanent residences.

Figure 22 Summary of Waterfront Regulated Land Use				
	Lake Area	Waterfront	% of	
	(Km²)	(Metres)	Total	
Miskwabi	2.68	10,782	100%	
WR4		5,199	48%	
WR4 Special		4,990	46%	
Open Space		35	0%	
Rural		558	5%	
Long	0.882	10,621	100%	
WR4		2,522	24%	
WR4L		4,268	40%	
Open Space		598	6%	
Rural		3,233	30%	
Wenona	0.591	5,500	100%	
WR4		1,448	26%	
WR4L		2,238	41%	
WR3L		362	7%	
Open Space		609	11%	
Rural		843	15%	
Negaunee	0.169	2,140	100%	
WR4		692	32%	
WR4L		245	11%	
Rural		1,203	56%	
Lakes Total	4,321	29,043	100%	
WR4		9,861	34%	
WR4L		6,751	23%	
WR4 Special		4,990	17%	
WR3L		362	1%	
Open Space		1,242	4%	
Rural		5,837	20%	

#### Zoning Legend:

WR4 – Waterfront Residential Type 4: residential uses; 45 m lot frontage; 20 m water building setback on lots developed as of 2004

WR4L – Waterfront Residential Type 4L: residential uses; 45 m lot frontage; 20 m water building setback on lots developed as of 2004; limited service (no public road)

WR4 Special – Waterfront Residential Type 4 Special: residential uses; WR4 with an additional building setback

WR3L – Residential Type 3L: residential uses; 45 m lot frontage;30 m water building setback; limited services (no public road)

**Open Space/Environmental** – no construction or use of any structure except for a private or public park

**Rural** – 20 ha lot area with a limited range of uses; permitted uses include a singlefamily dwelling, a bed & breakfast establishment, a hobby farm, a home business. Using Municipal Property Assessment Corporation (MPAC) assessment and parcel register data, a total of 451 privately-owned shoreline properties were identified on Long, Miskwabi, Wenona and Negaunee Lakes. Of those, only 66 or 14.6% were assessed as vacant.

	<b>Building Density on Shoreline Properties</b>					
	with Dwellings	Vacant Properties	Open Space or Shared Access	Total		
Miskwabi	142	21	12	175		
Long	118	24	0	142		
Wenona	89	14	3	106		
Negaunee	20	7	1	28		
Total of 4 Lakes	369	66	16	451		
Notes: based on analysis of MPAC assessment and register data in the spring of 2014.						

#### Figure 23 Building Density on Shoreline properties

There are also a large number of non-waterfront (back-lot cottages, rural and residential) properties within our watershed area. Using MPAC assessment and parcel register data, a total of 169 nonwaterfront properties were identified. Of these, 96 (57%) have dwellings and the remaining 73 are vacant (43%). Of the back-lot properties, 104 were identified as having deeded access to lakes, via shared ownership of the deeded-access lots detailed in the lake summaries above (66 with access to Miskwabi; 30 with access to Wenona; 8 with access to Negaunee).

Based on the foregoing discussion, there is still a significant number (143) of approved properties without buildings that can be developed on and around our lakes.

### LAND USE MAP



## **RESIDENTIAL OCCUPANCY**

Occupancy refers to the number of residential users and the duration of their stay at the lake. The number of people living on or using the lake has a direct impact on water quality. More cottages, longer visits, as well as conversion of seasonal to permanent residences will increase the amount of phosphorus generated from human waste and sewage as well as increased incidence of swimming and washing in the lakes. There will also be more contaminants entering the lakes as a result of increased use of fertilizers, more boating, increases in the number of shoreline trees being cut down and from more water runoff being channeled directly to our lakes. In addition, there is a growing concern about littering, noise and general mistreatment of the environment from increased occupancy. Development of water quality models often calculate estimates of phosphorus loading based on the total number of shoreline lots and whether the lot is used permanently or seasonally. They also consider information on the amount of time people stay on the lake as well as the estimated household size. More detailed information on residential occupancy will help to provide more accurate predictions of future water quality. The property owners' survey indicated that 79% of property owners are Seasonal versus 21% that indicated Full-Time occupancy. However, many property owners have reported that they are considering moving to the lake permanently and thus Full-Time occupancy could reach 38% in the future. The Average Number of Individuals in the Household was reported at 3.7 persons, considerably higher than the provincial average of 2.6 persons. Property owners use their property an average of 130 days a year, while54% use it in the winter. Approximately 8% of owners rent their dwelling and it was reported that this rate could increase to approximately 12% in the future.

# SEPTIC SYSTEMS AND WASTE DISPOSAL

E-coli bacteria from untreated human waste can represent serious health threats to humans. Phosphorus reaching the lake from malfunctioning sewage disposal systems will also speed lake eutrophication and threaten cold water fish (lake trout) populations and other inhabitants.

While there are a number of methods of treating waste water, by far the most common in our region is through a septic tank and leaching bed.<sup>42</sup> Although the Province sets the building code for septic disposal systems and takes care of licensing new technologies, permitting and inspections are handled locally. Up to now the Haliburton Kawartha Pine Ridge Health Unit has been contracted by the Municipality to enforce Part 8 of the Ontario Building Code. Beginning in July 2014 this responsibility changed to the Municipality.

Since 1973 all building permits in the Haliburton area require a comprehensive inspection and compliance with waste water system regulations prior to occupancy. As such, and given the rapid residential development since that time, the vast majority of properties on our lakes would have waste water treatment systems that at least at one time were compliant with government regulations. Septic system inspections are becoming commonplace in Real Estate transactions, which have the effect of revealing and remedying septic system deficiencies as properties are sold. In addition, the soil of the Canadian Shield surrounding the Haliburton area is rich with iron that acts as a natural buffer to phosphorus emissions. This further helps mitigate the risk of lake contamination.

<sup>&</sup>lt;sup>42</sup> Other methods include composting toilets, outhouses and holding tanks.

Nevertheless, there are certainly some older properties that would contain older waste water treatment systems.<sup>43</sup> Research from other similar cottage jurisdictions has also demonstrated that a relatively significant portion of septic systems required some form of remedial action.<sup>44</sup>

A malfunctioning septic system is one that allows untreated sewage effluent to enter the lake. This could be a result of:

- a damaged or leaking old tank
- a tank that has NOT been pumped in so many years that it allows solids to enter and clog the tile bed
- all soil pore space in a tile field becoming "clogged" with effluent minerals, and the effluent pools on the surface of the bed and runs overland into the lake
- the tile bed and perhaps underlying pipes/tiles have been damaged by being driven on.

There is no formal programme administrated by either the Health Unit or the Municipality to re-inspect septic systems. The onus rests with property owners to manage and maintain their septic systems and to ensure that the remnants of untreated human waste are not entering the lake water.

*Caring for your Septic System in Haliburton*, published by the Muskoka Watershed Council, lists a variety of practices property owners may undertake to help care for their systems:

- Obtain and review the permit for your septic system.
- Keep accurate records of pumping, maintenance and repair.

- Locate and prepare a sketch of components. Ensure all fixture drains are connected (as per design).
- Ensure all fixture drains are connected (as per design).
- Conserve water flowing to the system (i.e. low flush toilets).
- Repair any leaking plumbing fixtures.
- Do not put grease, paint or petroleum products down the drain.
- Flush only biodegradable products like toilet tissue; not dental floss, sanitary napkins, flushable wipes, cigarette butts, or paper towels.
- Avoid high water flows that may overload your system.
- Install an effluent filter to the outflow pipe leading from your septic tank to your leaching bed. An effluent filter will prevent suspended solids from entering the leaching bed and must be cleaned as prescribed by the manufacturer.
- Maintain vegetation on top of your leaching bed to aid in evaporation and prevent erosion; but do not allow trees or shrubs to grow too close to the bed or tank as their roots can clog or damage your system.
- Ensure renters/guests are aware of your system's proper use.

ISSUE: Malfunctioning septic systems are considered one of the most significant threats to the water quality of our lakes. This fact is all the more important for our lakes because most septic systems on the lakes, excluding those on the north shore of Miskwabi Lake, were installed over 30 years ago. ACTION:

- Support and work with the CHA education programme regarding pumping, inspection and maintenance of septic systems.
- Recommend that the Municipality review possible alternatives to prevent the renting of a cottage to more occupants than the septic system is designed to accommodate.
- Recommend that MACA work with the Municipality to undertake a Septic Re-Inspection Programme.

<sup>&</sup>lt;sup>43</sup> According to the survey, over 16% of property owners have owned their property for more than 40 years. As well, research from other cottage municipalities indicated that almost a third of septic systems are outdated.

<sup>&</sup>lt;sup>44</sup> Algonquin Highlands municipality (immediately northwest of Dysart et al) completed a 3-year-long re-inspection of all septic systems within their boundaries and found that 14% of approximately 5000 systems needed some form of remedial action.

## LAND USE REGULATIONS

Land-use planning in Ontario is regulated by the Planning Act of 1990 (as amended) under the Ministry of Municipal Affairs.<sup>45</sup>However, regional and local issues are dealt with at the county and municipal level. In our territory, the County Official Plan sets the framework for the Municipal Official Plan, out of which flow the regulatory zoning by-laws.

The purpose of the Official Plan is to establish a broad planning policy framework for the County, and is intended to guide the official plans and development approvals of the local municipalities. The Official Plan for the Municipality of Dysart et al was approved by the Minister of Municipal Affairs and Housing, March 11, 2004 and the text and schedules were updated December, 2010. <u>http://www.dysartetal.ca/PDFs/DysartOPConsolidationDec2010.pdf</u> As a consequence of the Official Plan, the Municipality updated its zoning by-laws

<u>http://www.emmersonlumber.com/pdf%20files/dysart\_bylaw.pdf</u> The scope of this document, which gets updated periodically, is to define the land uses through well-articulated zoning provisions, requirements for building permits and compliance. They stipulate the type and number of buildings that can be constructed on the lot and where they can be situated on the lot. Except for on the north shore of Miskwabi Lake, the minimum setback for building structures is 20 metres and 30 metres for leaching beds. Miskwabi Lake North Shore lots have an extra setback, varying by lot, and averaging between 20 to 40 metres.

Through enforcement of the zoning regulations the Municipality of Dysart et al has been successful in preventing the construction of any new boat houses on our lakes.

While there is a very specific process that needs to be followed to have zoning regulations changed or lot specifications split, changes do occur from time to time.

#### ISSUE:

Non-conforming land use or development on the lands adjacent to our lakes could threaten water quality, natural landscapes, and peace and quiet. This is particularly important as the lakes are a headwater area.

#### ACTION:

Work with the Municipality to reinforce all zoning and environmental regulations. Ensure MACA is informed of all potential re-zoning applications and has the opportunity to engage in the re-zoning assessment process.

<sup>&</sup>lt;sup>45</sup> In early 2014 the Province of Ontario released a revised 56-page Provincial Policy Statement (PPS) which provides direction for future land use planning. The PPS includes enhanced policy direction to better reflect the needs and unique circumstances of northern and rural Ontario including ensuring consideration of environmental lake capacity. The PPS 2014 became effective on April 30, 2014.

In addition, FOCA represents the interests of waterfront property owners relating to land use concerns. This association is lobbying to improve land use regulations so they better address waterfront specific issues (e.g. managing docks, in-water construction, building setbacks, septic systems, etc).

MACA has and continues to support these stewardship efforts.

# OTHER REGULATIONS AND GOVERNANCE

Waterfront properties are subject to a number of other legislative controls enforced at the federal, provincial and municipal levels. The federal Department of Fisheries and Oceans (DFO) enforces the Fisheries Act and the Navigable Waters Protection Act. The Fisheries Act makes it illegal for a person to do anything that harms fish habitat, or to deposit a harmful substance in waters containing fish. The Act provides for significant fines to be levied for infractions. The Ontario Ministry of the Environment regulates water quality through its Environmental Protection Act, which addresses public health concerns resulting from water contamination. Under the Act, spills must be reported, stopped and controlled. The Ministry is also responsible for the approval of waste disposal systems greater than 10,000 litres. The Ministry also oversees and grants permits for substantial water extractions.

The Ministry of Natural Resources regulates the province's natural resources in lake areas in accordance with the Public Lands Act and the Lakes and Rivers Improvement Act. These Acts require a work permit before work around water bodies can occur, including dredging; filling; blasting; removal of vegetation; and construction of retaining walls, docks and boathouses.



Miskwabi Clearing

Original Artwork: Al Van Mil

# **GOING FORWARD**

Through the four-year process of developing the Miskwabi Area Watershed Plan the people of the community have clearly revealed their values, as well as the lifestyle they enjoy and the lake characteristics that they believe are essential. Understanding these factors has led to an action plan of stewardship and regulatory recommendations.

The individuals of the community vary substantially economically, in interests and activities: however, all agree that it is essential to maintain water quality and natural landscapes which are the body and soul of the community.

The Miskwabi Area Watershed Plan is a living document based on the input of the members of the community and intended for use by all. It is a beginning, not an end; a report not to be put on a shelf but to be used and updated on a regular basis. In this regard, at the July 5, 2014 MACA Annual General meeting the Plan was unanimously endorsed by the membership. As well, it was agreed that a Director's role on the Board will be established to guide the implementation of the Plan and its on-going relevance for the community.



Miskwabi Triptych

Original Artwork: Annette Blady

# **APPENDICES**

# **BIBLIOGRAPHY and REFERENCES**

Aquatic Research Section, MNR Ontario. Summary: Defining a Dissolved Oxygen Criterion for the Protection of Lake Trout Habitat accessed at www.mnr.gov.on.ca on Feb. 2, 2014. Balch, Gordon. Ecosystems Integrity: Methods in Environmental Science, Fleming College Dec. 2011. Balch, Gordon. Environmental Assessment concepts, framework and policy. Fleming College Dec. 2011. Biodiversity Branch, Ministry of Natural Resources. 2006. Dissolved Oxygen Criterion for the Protection of Lake Trout Habitat. accessed at www.mnr.gov.on.ca on Feb. 2, 2014. Bird Studies Canada. The Canadian Lakes Loon Survey 1981-2012 Bosley, Holly E. 2005. Techniques for Estimating Boating Carrying Capacity: A Literature Review Cadman, M.D. et al. 1987 Ontario Breeding Bird Atlas 2000 – 2005. Waterloo: Federation of Ontario Naturalists. Central Lake Ontario Conservation Authority Long-Term Aquatic Monitoring Program 2009 - 2013 Coalition of Haliburton Property Owners' Association 2011. Living on the Edge, A Handbook for Lake Stewards. County of Haliburton Official Plan accessed at www.haliburtoncounty.ca/services/planning-and-gis/official-plan-2. County of Haliburton GIS & Mapping Site. www.haliburtoncounty.ca/services/planning-and-gis County of Haliburton, Moose and Deer Yards. accessed at www.haliburton.ca/wp-content/uph/2013/10/OPSscheduleA.pdf. Department of Lands and Forests, Ontario. Report of Preliminary Survey-Atwell's Lake, memorandum 1958. Doshi, Sheela. 2006. Recreational Carrying Capacity in Lakes: How much is too much? Ducks Unlimited. 2011. Enhanced Wetland Classification – Inferred Products User Guide. Version 1.0 Ducks Unlimited. 2004. The Importance of Wetlands and Upland Conservation Practices in Watershed Management. Farrar, John Laird. Trees in Canada. 2011 Filzhenry & Whiteside Limited Fisheries and Oceans Canada. 2008. The Shore Primer, Ontario Edition. Federation of Ontario Cottagers Associations (FOCA). Lakeshore Capacity Assessment: The Facts Geological Highway Map Southern Ontario, revised 1979 published by Ministry of Natural Resources (includes Physiographic Map) Haliburton Highlands Lands Trust. Natural Heritage Planning Framework: Review of the setup process for Natural Heritage Systems. Haliburton Highlands Land Trust. 2007. County of Haliburton Natural Heritage Mapping: A Compilation and Preliminary Assessment prepared by Glenside Consulting. Haliburton Highlands Land Trust. 2008 Species at Risk Natural Heritage Strategy prepared by Paul C. Heaven accessed Sept. 27, 2011

Hyatt, Crystal. 2020. Lakeshore Capacity assessment: The Facts. Federation of Ontario Cottagers Associations (FOCA).

Idika, Odum. 2010. A preliminary investigation into the ecological significance of headwater drainage features in Southern Ontario. University of Waterloo.

Knopf, Alfred A. 1979 The Audubon Society Field Guide to North American Wildflowers. Chanticleer Press Edition.

Kusler, J. A. 1972. Carrying Capacity Controls for Recreation Water Users. Upper Great Lakes Regional Commission.

Lakeshore Capacity Assessment Handbook: protecting Water Quality in Inland lakes on Ontario's Precambrian Shield. 2010

Ontario MOE, MNR & MMAH accessed at http: www.ene.gov.on.ca/publications/7642e.pdf.

Lake Planning Handbook for Community Groups. FOCA and French Planning Services Inc.

Lake Ripley Watercraft Census & Recreational Carrying Capacity Analysis. 2003. Prepared by Lake Ripley Management District.

Lake Simcoe Water Quality Update. 2010. Ontario Environmental Monitoring and Reporting Branch.

MacCulloch, Ross Douglas. 2002. The ROM Field Guide to Amphibians and Reptiles in Ontario. Co-published by McClelland & Stewart Ltd. and the Royal Ontario Museum.

Ministry of Environment Ontario. 2011. Water Quality and Management of Lake Trout Lakes.

Ministry of Natural Resources, Ontario. www.mnr.on.ca.

Ministry of Natural Resources, Ontario. 2005. Summary: defining a dissolved oxygen criterion for the protection of lake trout habitat.

Ministry of Natural Resources, Ontario. 2006. Dissolved Oxygen Criterion for the Protection of Lake Trout Habitat. Fisheries Policy 7.5.1

Ministry of Natural Resources, Ontario. 1990. Environmental guidelines for Access Roads and Water Crossings.

Ministry of Natural Resources, Dr. David Evans, Haliburton Lake Trout Adoptive Management Study, accessed at

www.mnr.gov.on.ca/en/Business/Aquatics on June 16, 2013

Ministry of Natural Resources. W. Wilson. Miskwabi Lake Fisheries Report 1980.

Ministry of Natural Resources, Ontario. *Moose Resource Report*. accessed at <u>www.ontario.ca/moose</u> on Feb. 17, 2014.

Ministry of Natural Resources, Ontario. Ontario Wetland Evaluation System, Southern Manual.

Ministry of Natural Resources, Ontario. Significant Wetlands and the Ontario Wetland Evaluation System. www.mnr.gov.on.ca

Ministry of Natural Resources. Species at Risk. www.mnr.on.ca accessed on June 24, 2014.

Ministry of Natural Resources. What are Wetlands?

http://www.mnr.gov.on.ca/stdprodconsume/groups/lr/@mnr/@biodiversity/documents/document/stdprod\_091595.pdf. Municipality of Dysart et al. *Official Plan* accessed @ www.dysartetal.ca/wp-content/uploads/2013/09

National Wetlands Working Group. 1997. www.wetlandpolicy.ca

Ontario Invasive Plant Council. Grow Me Instead. accessed @ www.ontarioinvasiveplants.ca

Ontario Woodlot Association. 2009. A Landowner's Guide to Careful Logging.

#### Popple, Robert.2003. Northern Belle, The Life of Haliburton's Ethel Curry. RTP Publications.

Quinlan, R., A.M. Paterson, J.P. Smol, M.S.V. Douglas, and B.J. Clark. 2005. Comparing different methods of calculating hypolimnetic oxygen

(VWHO) in lakes. accessed at Dorset Environmental Science Centre www.desc.ca Feb 2014.

SFI Implementation Committee Ontario. A Guide to Best Management Practices for Forest Operations in Northern Ontario.

Shoreline Owner's Guide to Lakeland Living. Lakeland Alliance

State of Ohio, Environmental Protection Agency. The Importance and Benefits of Primary Headwater Streams

www.epa.ohio.gov/portals/35/wgs/headwatersHWH import jan2003.pdf

Stroud Water Research Center. Protecting Headwaters: The Scientific Basis for Safeguarding Stream and River Ecosystems.

http://www.stroudcenter.org/research/PDF/ProtectingHeadwaters.pdf

*The Wolves of Algonquin Park.* 2000. Report by the Algonquin Wolf Advisory Group.

Toronto and Region Conservation. Headwater Study 2013

Toronto Regional Conservation Authority. 2007. The Natural features of Headwater Drainage Features: A literature review. www. trca.on.ca

# ACRONYMS and ABBREVIATIONS

ABBO	Atlas of the Breeding Birds of Ontario
СНА	Coalition of Haliburton Property Owners Associations
DFO	Department of Fisheries and Oceans, Canada
EDDMapS	Early Detection & Distribution Mapping System
GIS	Global Information Systems
MACA	Miskwabi Area Community Association
	formerly Miskwabi Area Cottagers' Association
MNR	Ontario Ministry of Natural Resources
MPAC	Municipal Property Assessment Corporation
MOE	Ministry of the Environment, Ontario
OFAH	Ontario Federation of Anglers and Hunters
OPP	Ontario Provincial Police
PSW	Provincially Significant Wetland
SARO	Species at Risk Ontario
TSW	Trent-Severn Waterways



Miskwabi Wolf Original Artwork: Gladys Gottschalk

# LIST OF MAPS

1. Topographic Map, Miskwabi Area Watershed	pg.17
2. Satellite Image of Miskwabi Area	18
<ol><li>Miskwabi Area Water Flow Map</li></ol>	19
4. Physiographic Map of Southern Ontario	20
5. Map 2441 Ontario Geologic Survey	20
6. Bathymetry Map, Long Lake	22
7. Bathymetry Map, Miskwabi Lake	23
8. Bathymetry Map, Negaunee Lake	24
9. Bathymetry Map, Wenona Lake	25
10. Miskwabi Area Roads	28
11. Miskwabi Area Catchment Area	37
12. Miskwabi Area Wetlands	39
13. Miskwabi Area Streams	41
14. Moose Late Wintering Area	47
15. Land Use Map	62



### Trapper's Trail Clearing

Original Artwork: Al Van Mil

# LIST OF FIGURES

1. Stakeholder Survey Results, Natural Features	pg.4
2. Stakeholder Survey Results, Issues	4
3. Stakeholder Survey Results, Land Activities	5
4. Stakeholder Survey Results, Water Activities	5
5. Developer's Sales Brochure	15
6. MACA original logo	16
7. MACA logo adopted in 2012	16
8 Miskwabi/Long Lakes Historical Water Levels	26
9 Miskwabi/Long Lakes Preferred Water Levels	27
10. Water Quality Parameters	30
11. Shoreland Zones	35
12. List of Miskwabi Area Wetlands	38
13. List of Miskwabi Area Permanent Streams	40
14. List of Haliburton Mammals	46
15. Seasonal Movements of Moose in Ontario	47
16. Uncommon Birds of Miskwabi Area	48
17. ABBO Breeding Evidence in Miskwabi Area	50
18. List of Haliburton Herpetofaunal	51
19. Miskwabi Area Species at Risk	52
20. Comparison of Lake Surface Area and Perimeter	58
21. Recreational Boat Carrying Capacity Table	59
22. Summary of Water Front Regulated land Use	61
23. Building Density on Shoreline Properties	61
#### ACKNOWLEDGEMENTS

The Miskwabi Area Watershed Plan Committee acknowledges the financial support of the members of the Miskwabi Area Community Association (MACA).

#### Invaluable input was provided by:

David Flowers, Fisheries Extension Biologist, Ontario Ministry of Natural Resources, Minden Pat Martin, Planning Director, Municipality of Dysart et al

#### Professional Consultation:

Randy French, French Planning Services, Inc.

#### Special Thanks to:

Don Benson, Mountain Lake Property Owners Association Andy Mosher, Research and assistance Jane Tousaw, Director of Planning, County of Haliburton Nicole Tuyten, formerly Stewardship coordinator, Haliburton Highlands Stewardship Council Dr. Brent Wooten, Director Applied Research, Fleming College

#### **Original Artwork:**

The natural beauty of the Miskwabi lakes area inspires artists of many mediums. Our thanks to the following Miskwabi area artists for permission to reproduce their work here and on the Introductory DVD.

Annette Blady
<b>Gladys Gottschalk</b>
Susan Hay

Tim Payment Al Van Mil

#### Printing:

Printed by Autumn Marketing of Haliburton County.

1<sup>st</sup> printing August 2014 2<sup>nd</sup> printing January 2015

#### Photographs:

Special thanks to the following Miskwabi area photographers whose local photos enliven the printed version of the Watershed Plan as well as the Introductory DVD.

Terry Anderson William Bunn Michael Giza Larry Holden Susanne James Dave McCulloch Ken Morrison Andy Mosher Tim Payment Tom Pendlebury Mike Tipping Virginia Vranckx Phil Wiseman

#### Introductory DVD:

The DVD was conceived, written and recorded by Rick Despard in his Miskwabi Lake studio.



Wenona Lake

Original Artwork: Susan Ha

# **An Integral Vision**

### For today and the future:

Pristine headwater lakes Natural landscapes Healthy environment Active lifestyles

Nothing is separate. Everything in the watershed is interconnected.





Peter Dilworth Long



**Rick Despard** Miskwabi



Michael Giza Miskwabi



Larry Holden Miskwabi



Miskwabi



# **MISKWABI AREA WATERSHED PLAN**

## **TEAM**

Camera shy: Stuart Buck, Miskwabi Haden Heathcock, Miskwabi



Phyllis McCulloch Wenona



**Tim Payment** Tom Pendlebury Miskwabi



Neil Roberts Long

Virginia Vranckx





**STAKEHOLDER WORKSHOPS** 

August 31, 2013 and June 29, 2014